

Classification of Singularities and Bifurcations of Critical Points of Even Functions

E.A. Kudryavtseva*, E.L. Lakshtanov †

Abstract

Singularities of even smooth functions are studied. A classification of singular points which appear in typical parametric families of even functions with at most five parameters is given. Bifurcations of singular points near a caustic value of the parameter are also studied. A determinant for singularity types and conditions for versal deformations are given in terms of partial derivatives (not requiring a preliminary reduction to a canonical form).

1 Introduction

In this work, we consider families of smooth even functions $\omega_\lambda(k) : \mathbb{T}^\nu \rightarrow \mathbb{R}$ on a torus $\mathbb{T}^\nu = (S^1)^\nu$ where $S^1 = [0, 2\pi]/0 \sim 2\pi$ and the parameter λ takes values in some domain $\mathbb{R}^l$:

$$\omega_\lambda(k) = \omega_\lambda(-k), \quad k \in \mathbb{T}^\nu, \quad \lambda \in \mathbb{R}^l. \quad (1.1)$$

We will study bifurcations of critical points for smooth families of even smooth functions, their singularities, and, in particular, obtain a classification of degenerate critical points for generic families of germs of even functions, which are determined in a neighbourhood of zero (or in a neighbourhood of a fixed point of the involution $k \mapsto -k$), with the number of parameters $l \leq 5$. Moreover, we will show how to determine types of critical points of even functions which appear in typical parametric families with $l \leq 4$ parameters, and formulate conditions on their deformations to be typical (more precisely, versal), in terms of partial derivatives (not requiring a preliminary reduction to a canonical form).

The study of properties of families of even functions appears in problems about planetary systems with satellites, integrable Hamiltonian systems with 2 degrees of freedom, problems of spectral analysis of stochastic operators of multiparticle systems, and problems involving two (quasi)particles interaction. It was shown in [5, 6, 7, 8] how the type of singularities of a family $\omega_\lambda(k)$ affect the spectrum structure. An extremely important task is therefore to describe *stable families* of even functions in a small neighbourhood of a singular point (they are also called *families in generic position* or *typical families*), i.e. families which do not change their properties under small perturbations (for the precise definition see 4.9).

We tried to make this work readable for those readers who are not familiar with singularity theory, as well as to demonstrate, using simple examples, how to use results of [1] which are very important for applications.

The authors are grateful to S.M. Gusein-Zade, M.E. Kazaryan, G. Wassermann, and V.M. Zakalyukin for useful discussions.

*Moscow State University. e-mail: eakudr@mech.math.msu.su

†Moscow State University. e-mail: lakshtanov@rambler.ru

1.2 Critical points of even functions on a torus $\mathbb{T}^\nu$

The automorphism $k \mapsto -k$ of the torus has 2^ν fixed points

$$(k_1, \dots, k_\nu), \quad k_i \in \{0, \pi\} \subset S^1, \quad i = 1, \dots, \nu.$$

These points are obviously critical. We will call them *basic*. Observe that, if a non-basic point $k \in \mathbb{T}^\nu$ is critical, then the point $-k$ is also critical. Thus, non-basic critical points appear in pairs $(k, -k)$, moreover the degeneracy type of these points is the same. We will call such points *twin* or *additional*.

Recall [1, v. 2, 6.1.Γ] that a value of the parameter λ is called *caustic* if the function $\omega_\lambda(k)$ admits a degenerate critical point. It is obvious that, if some value $\lambda = \lambda_0$ is not caustic, the number of critical points and their types remain the same in a neighbourhood of λ_0 . Thus, appearance and disappearance of critical points is related with the passage of the parameter λ across hypersurfaces (i.e. surfaces of codimension one) of caustic values. For a detailed study of these processes, let us describe the types of degenerate critical points, which are non-removable for the entire family of functions. In particular, this is necessary for study of oscillating integral asymptotics, see below.

2 Classification of singularities of even smooth functions

Thus, for even functions $\omega_\lambda(k)$, there are two types of critical points. In a neighbourhood of a basic critical point, the function $\omega_\lambda(k)$ is even, while, in a neighbourhood of an additional critical point, it is an arbitrary function (i.e. a function which is not assumed to be symmetric relative to the critical point). In the latter case, we can use the classification from [1].

2.1 A survey of the classification of critical points of arbitrary functions due to [1, v. 1, 11.2]

We present here a table of germs of smooth functions at zero up to the following transformations: adding a constant, adding a nondegenerate quadratic form in the remaining variables (the number of these variables equals the rank of the second differential of the function at zero), and a smooth change of variables leaving the origin fixed. The integer c in the table is the *codimension of the class of singularities* in the space of function germs at zero. It equals the minimal number $l \geq c$ of parameters of a family, such that a critical point of the class under consideration is non-removable under small perturbations of the family. The integer μ is the *multiplicity* of the critical point 0 (see [1, v. 1, 6.3] and 3.3). For all known types of singularities, it equals the *codimension* of the singularity in the space of function germs at zero plus 1 (see 4.9 and 4.10).

Remark 2.2 Recall the definitions of the notions *singularity* and *class of singularities* from [1] mentioned above. If two functions can be obtained from the same function by the above mentioned transformations, one says that these functions have the same *type of critical point* or the same *type of singularity* at zero, and the germs of these functions at zero are R^+ -equivalent (more precisely, stably R^+ -equivalent [1, v. 1, II]). The notion of R -equivalence is similarly determined, where adding a constant is forbidden, see [1] and 3.4. An R^+ -equivalence class of germs, i.e. the space of all germs with a given type of singularity, is also called a *singularity*. Singularities which can be joined to each other by a smooth path in the space of singularities having the same multiplicity μ form a *class of singularities*, more precisely a *class of μ -equivalent singularities*, or a $\mu = \text{const}$ *stratum*, see 4.10 and [1, v. 1, 15.0.1]. A class of singularities is (at least for all known finite-multiple singularities) a smooth parametric family of singularities. For such a parametric family, the parameters are called *moduli*, with the number m of parameters being called the *modality* of

the singularity (for a more general definition of modality see the introduction of [1, v. 1, II]). The equality $\mu = c + m + 1$ is valid, where m is the modality and c is the codimension of the class of singularities, see 4.10 and [1, v. 1, 15.0.4]. See also the examples after Remark 4.10.

We remark that the singularities listed in Table (2.3) do not have moduli ($m = 0$), thus they are *simple* [1, v. 1, II]. For generic families of functions with $l \leq 5$ parameters, there are no critical points apart from those which are equivalent to the singularities listed in Table (2.3).

Type of singularity	Normal form	Restrictions	μ	c	β
$A_k^\pm$	$\pm x^{k+1}$	$k \geq 1$	k	$k - 1$	$\frac{k-1}{2k+2}$
$D_k^\pm$	$x^2 y \pm y^{k-1}$	$k \geq 4$	k	$k - 1$	$\frac{k-2}{2k-2}$
$E_6^\pm$	$x^3 \pm y^4$	—	6	5	$\frac{5}{12}$

(2.3)

Seven of these singularities, with $c \leq 4$, are known in catastrophe theory as the “seven of Thom”. Unimodular singularities ($m = 1$) appear for $c = 6$, however we do not consider them here.

Remark 2.4 The integer β is the *singularity exponent* of a given critical point. It can be determined by means of the asymptotics of the oscillating integral

$$\int_{U(0)} e^{itf(x)} \varphi(x) dx_1 \dots dx_\nu \sim \text{const} \cdot t^{\beta-\nu/2}, \quad t \rightarrow \infty.$$

Here, the integration is taken over a small neighbourhood of the point $x = 0 \in \mathbb{R}^\nu$, which is a critical point of the phase $f(x)$ with singularity type under consideration, and the amplitude $\varphi(x)$ does not vanish in $U(0)$. (For more details see [1, v. 2, p. 134].)

Remark 2.5 The listed types of critical points never correspond to a (local) extremum (i.e. a point of local minimum or local maximum) apart from the types A_{2k-1} , $k \geq 1$. Moreover, Table (2.3) implies that for generic one-parameter families, an additional extreme point never degenerates. V.A. Vasiliev composed a table of germs of smooth functions in a neighbourhood of minimum points, which occur in generic families of functions with $l \leq 16$ parameters, see [1, v. 1, 17.2].

2.6 Classification of critical points of even functions

We present here a table of germs of smooth even functions at zero up to the following transformations: adding a constant, adding a nondegenerate quadratic form in the remaining variables, and a smooth odd change of variables. The integer c_e in the table is the *even codimension of an even class of singularities* and equals the minimal number $l \geq c_e$ of parameters of a family, such that a critical point of the class under consideration is non-removable under small perturbations of the family of even functions. The integer μ_e is the *even multiplicity* of the function at zero (see 3.3). It equals the even codimension of the singularity plus 1.

The notions *R_O -equivalence* of even germs (see 3.4), *even singularity*, *even class of singularities*, *even type* of a critical point, *even modality* m_e , and *even moduli* are defined similarly to 2.2. The equality $\mu_e = c_e + m_e + 1$ is valid.

Even singularities of the series $A_{e,k}$ do not have moduli ($m_e = 0$). Other even singularities listed in Table (2.7) are unimodal ($m_e = 1$). Actually, all finite-even-multiple even singularities but $A_{e,k}$ have moduli ($m_e \geq 1$) [12, Theorem 4.3]. For generic families of even functions with $l \leq 5$ parameters, there are no even singularities at zero apart from those which are equivalent to the even singularities listed in Table (2.7).

Even class of singularities	Normal form	Restrictions	μ_e	c_e	β
$A_{e,k}^\pm \subset A_{2k-1}^\pm$	$\pm x^{2k}$	$k \geq 1$	k	$k - 1$	$\frac{k-1}{2k}$
$X_{e,5}^{\pm\pm} \subset X_9^{\pm\pm}$	$\pm x^4 + ax^2y^2 \pm y^4$	$a^2 \neq 4$, if $++$ or $--$	5	3	1/2
$X_{e,r+3}^{\pm\pm} \subset X_{2r+5}^{\pm\pm}$	$\pm x^4 \pm x^2y^2 + ay^{2r}$	$a \neq 0$, $r \geq 3$	$r + 3$	$r + 1$	1/2
$Y_{e,r,s}^{\pm\pm} \subset Y_{2r,2s}^{\pm\pm}$	$\pm x^{2r} + ax^2y^2 \pm y^{2s}$	$a \neq 0$, $r, s \geq 3$	$r + s + 1$	$r + s - 1$	1/2
$\tilde{Y}_{e,r}^\pm \subset \tilde{Y}_{2r}^\pm$	$\pm(x^2 + y^2)^2 + ay^{2r}$	$a > 0$, $r \geq 3$	$2r + 1$	$2r - 1$	1/2
$Z_{e,7}^\pm \subset Z_{13}^\pm$	$x^3y \pm y^6 + axy^5$	—	7	5	5/9

(2.7)

Table (2.7) is an extension of the table of even germs of even multiplicity $\mu_e \leq 5$ from the diploma work of M. Beer, see [4, Satz 5.6] or [12, Theorem 4.1].

Remark 2.8 A singularity $X_{e,5}$ is a (local) minimum only if its normal form is $x^4 + ax^2y^2 + y^4$ where $a > -2$, $a \neq 2$. Singularities $X_{e,r+3}$ and $Y_{e,r,s}$ are (local) minima only if the normal form is $x^4 + x^2y^2 + ay^{2r}$, or resp. $x^{2r} + ax^2y^2 + y^{2s}$ with $a > 0$. A singularity $\tilde{Y}_{e,r}$ is a (local) minimum only if the normal form is $(x^2 + y^2)^2 + ay^{2r}$.

The result formulated above is proved in Section 3. Section 4 gives the definition and properties of even versal deformations of germs, and explains their relation to generic parametric families of smooth even functions. The reader who is mainly interested in a description of bifurcations of critical points for generic families of smooth even functions can pass to Section 5.

3 Germs of even functions

3.1 Equivalence of germs

Let E be the algebra of germs at zero of C^∞ -functions $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$, and let $\mathfrak{m} \subset E$ be the subalgebra of germs of functions, whose value at zero vanishes. Further $E_e \subset E$ is the subalgebra of germs of even functions, and $\mathfrak{m}_e = \mathfrak{m} \cap E_e$ is the maximal ideal of the subalgebra E_e .

Let $f \in \mathfrak{m}_e$. Then all partial derivatives $\frac{\partial f}{\partial x_i}$ are odd functions. Observe that linear combinations of the form $h_1 \frac{\partial f}{\partial x_1} + \dots + h_\nu \frac{\partial f}{\partial x_\nu}$ with coefficients $h_k \in E$, which are odd functions, form an ideal $I_{\nabla f}^e \subset \mathfrak{m}_e$ of the algebra E_e . This ideal is called the *even gradient ideal*, or the *even Jacobian ideal*, of the germ f .

Definition 3.2 The *even local algebra of the gradient map of f at zero* is the quotient algebra of the algebra of even germs, having a vanishing value at zero, by the ideal which is generated by the components of ∇f :

$$Q_{\nabla f}^e = \mathfrak{m}_e / I_{\nabla f}^e.$$

Example. Let $\nu = 1$ and $f(x) = x^6$. Then $Q_{\nabla f}^e = \{a_2x^2 + a_3x^4\}$. (The divided powers algebra with degrees less than 5 and a vanishing free term.)

Definition 3.3 The integer $\mu_e = \dim Q_{\nabla f}^e + 1$ is called the *even local multiplicity* of the gradient map of $f \in E_e$ at zero, or the *even multiplicity* of the critical point 0 of the function $f \in E_e$ (compare [1, v. 1, 6.3]). If $\mu_e < \infty$ then the critical point 0 of the even function f is called a *finite-even-multiple* critical point.

Let G be the group of germs of smooth odd changes of variables in $\mathbb{R}^\nu$, i.e. germs of diffeomorphisms h :

$$h(0) = 0, \quad h(x) = -h(-x).$$

The change of variables by means of $h \in G$ obviously transforms even functions to even ones, and odd functions to odd ones. Besides, under such a change of variables, the ideal $I_{\nabla f}^e$ and the algebra $Q_{\nabla f}^e$ are equivariantly transformed: $I_{\nabla(h^*f)}^e = h^*I_{\nabla f}^e$ and $Q_{\nabla(h^*f)}^e = h^*Q_{\nabla f}^e$. Therefore, the even multiplicity μ_e of the critical point 0 is preserved under odd changes of variables in $\mathbb{R}^\nu$.

Definition 3.4 Elements $f_1, f_2 \in E_e$ which admit a change of variables $h \in G$ such that

$$f_1 \circ h^{-1} \equiv f_2$$

are called R_O -equivalent, i.e. *right-odd equivalent*.

Recall that (not necessarily even) function germs f_1, f_2 are called R -equivalent if the equality $f_1 \circ h^{-1} \equiv f_2$ is valid for some, not necessarily odd, diffeomorphism h with $h(0) = 0$.

The following analogues of the Morse Lemma, the splitting lemma, and the Tougeron theorem about a finite-determination show that the class of odd changes of variables is wide enough:

Theorem 3.5 («Morse lemma» for even functions) *In a neighbourhood of the nondegenerate critical point 0, an even function is R_O -equivalent to the sum of a quadratic form and a constant.*

The above theorem is a partial case of the so called equivariant Morse lemma, see [1, v. 1, 17.3].

Theorem 3.6 (Splitting lemma for even functions) *In a neighbourhood of the critical point 0 of co-rank $k \leq \nu$, an even function is R_O -equivalent to a function having the form $\psi(x_1, \dots, x_k) + Q(x_{k+1}, \dots, x_\nu)$ where ψ is an even function in k variables, and Q is a nondegenerate quadratic form in $\nu - k$ variables.*

Theorem 3.6 generalizes Theorem 3.5. It is proved by means of the «parametric Morse lemma» for even functions. Theorem 3.6 is a partial case of the $\mathbb{Z}_2$ -splitting lemma [4], which is a partial case of the equivariant splitting lemma [12, Lemma 2.1]. For arbitrary smooth functions, a similar assertion is proved by J. Mather in his unpublished notes about R -equivalence. For the case $k = 1$, a proof is given in the beginning of the proof of the lemma from [1, v. 1, 9.6].

Theorem 3.7 (Tougeron theorem about finite-determination of an even germ) *In a neighbourhood of the finite-even-multiple critical point 0, an even function is R_O -equivalent to a polynomial (namely, to its Taylor polynomial of degree $2\mu_e$ at 0 where μ_e is the even multiplicity).*

Actually, this theorem admits the following generalization: one can choose the neighbourhood of 0 in such a way that any even function which is close enough to the function under consideration is also R_O -equivalent in this neighbourhood to a polynomial of degree $2\mu_e$ (namely, to its Taylor polynomial of degree $2\mu_e$ at 0).

A proof of this theorem (as well as the generalization from above) is analogous to the proof of the Tougeron theorem about finite-determination of a usual germ (for a proof by J. Mather see [1, v. 1, 6.3 and 6.4]), see [9] and [4].

Example. For a nondegenerate critical point ($\mu_e = 1$), an even function is R_O -equivalent to its Taylor polynomial of degree 2 («Morse lemma»). This example shows that the degree $2\mu_e$ can not be replaced by a lesser one.

3.8 Classification of singularities of even germs

In order to obtain a classification of singularities of even smooth functions, we can not unfortunately just consider all even normal forms of singularities of the usual smooth functions, since we do not know whether the R -equivalence of even germs with a vanishing value at 0 implies their R_O -equivalence. But we can transfer the corresponding classification assertions for the singularities of usual germs (see [1, chap. 2]) to our case.

Theorem 3.9 *The space of germs at 0 of even functions in ν variables with the critical point 0 of co-rank $k \leq \nu$ has codimension $k(k+1)/2$ in the space of germs of even functions at 0. (The co-rank of a critical point is the co-rank of the second differential of the function at this point.)*

In particular, in generic families of even functions with at most 5 parameters, only singularities of co-rank at most 2 occur. Thus, due to Theorem 3.6, in order to classify singularities and even singularities of functions which occur in generic families with at most 5 parameters, one may restrict oneself to functions in two variables.

A proof of the above theorem follows from the fact that the codimension of the set of matrices of co-rank k in the space of all symmetric matrices of order ν equals $k(k+1)/2$, see [1, v. 1, 2.2 and 11.1]. Here one should take into account that the second differential of an even function at zero can be an arbitrary symmetric matrix.

Theorem 3.10 *Let 0 be a finite-even-multiple critical point of co-rank 1 of an even function. Then, in a neighbourhood of 0, the function is R_O -equivalent to the function $c + x^{2\mu_e} + Q$ where c is a constant, μ_e is the even multiplicity, and Q is a nondegenerate quadratic form in the remaining variables.*

A proof is obtained by means of Theorem 3.6. It is analogous to the proof of the similar assertion for arbitrary smooth functions, see the lemma from [1, v. 1, 9.6] and Statement 6.3.

Thus, in order to describe all singularities in generic (i.e. stable, see 4.9) families of even functions with $l \leq 5$ parameters, it remains to classify even singularities of co-rank 2 of even functions in two variables. Let us show that any even singularity of interest has a nonvanishing 4-jet (i.e. its Taylor series contains a nonvanishing term of order 4 or less).

Theorem 3.11 *The space of germs at 0 of even functions in two variables with a vanishing 4-jet has codimension 8 in the space of germs of even functions with critical value 0 at the point 0.*

A proof of the above theorem follows from the invariance of the order of the lowest term under a diffeomorphic change of variables. Moreover, one should observe that the condition about the absence of the second order terms is defined by 3 equations, see Theorem 3.9, while the condition about the absence of the fourth order terms is defined by 5 equations.

3.12 Even singularities of co-rank 2 with vanishing 2-jet and nonvanishing 4-jet

Consider a real-valued 4-form in two variables

$$A(x, y) = a_1x^4 + a_2x^3y + a_3x^2y^2 + a_4xy^3 + a_5y^4, \quad a_i \in \mathbb{R},$$

where at least one of the coefficients a_i does not vanish. Zeros of the equation $A(x, y) = 0$ define four lines on the complex plane (x, y) . Thus the form A can be represented as the product of four linear forms $\tilde{A}_i = u_ix + w_iy$ with complex coefficients, $i = 1, 2, 3, 4$. Combining the linear forms into pairs, one can obtain a representation

$$A(x, y) = A_1(x, y)A_2(x, y)$$

where $A_i(x, y)$, $i = 1, 2$ are quadratic forms with real coefficients. We obtain the following cases of canonical forms of real-valued 4-forms in two variables.

1. If the four lines $\tilde{A}_i(x, y) = 0$ are pairwise different, or the lines of one of the pairs are not real and the lines of the other pair are different, then the quadratic forms A_i are nondegenerate. Moreover, they can be simultaneously reduced to diagonal form by a real linear change of variables (after a change of the grouping of the lines into pairs if necessary). This reduces the form $A(x, y)$ to the form

$$A(x, y) = \pm x^4 + ax^2y^2 \pm y^4, \quad a \in \mathbb{R}.$$

2. If exactly two of the real lines coincide, for example, $\tilde{A}_1(x, y) = 0 \Leftrightarrow \tilde{A}_2(x, y) = 0$, then one of the quadratic forms $A_i(x, y)$ is nondegenerate, and the second one is degenerate. Moreover, they can be simultaneously reduced to diagonal form by a real linear change of variables. This reduces the form $A(x, y)$ to the form

$$A(x, y) = x^2(\pm x^2 \pm y^2).$$

3. If the four lines are real and the lines of any pair coincide, while the lines of different pairs are different (after a change of the grouping of the lines into pairs if necessary), for example $\tilde{A}_1(x, y) = 0 \Leftrightarrow \tilde{A}_2(x, y) = 0$ and $\tilde{A}_3(x, y) = 0 \Leftrightarrow \tilde{A}_4(x, y) = 0$, then, after a real linear change of variables, the form $A(x, y)$ looks like

$$A(x, y) = \pm x^2y^2.$$

4. If exactly three (real) lines pairwise coincide then, after a real linear change of variables, the form $A(x, y)$ looks like

$$A(x, y) = x^3y.$$

5. If all four (real) lines coincide then, after a real linear change of variables, the form $A(x, y)$ obviously looks like

$$A(x, y) = \pm x^4.$$

We remark that all changes of variables from above are linear and therefore are *odd*.

Let us notice at once that the case 5 is not of interest to us, since the codimension of the subspace of even germs with a 4-jet (i.e. the Taylor polynomial of degree 4) of such type equals 6. Thus, such a singularity can not appear in generic families of even functions with $l \leq 5$ parameters. In fact, the subspace of such germs is defined by three equations on the quadratic terms and three equations on the fourth order terms.

Consider the case 1. If $|a| = 2$ and the signes $\pm\pm$ in the form $A(x, y)$ equal $++$ or $--$ then $A(x, y) = \pm(x^2 \pm y^2)^2$. Therefore, the form $A(x, y)$ has the form $\pm x^2y^2$ (case 3), or $\pm(x^2 + y^2)^2$ (the latter subcase will be considered after case 3).

Theorem 3.13 (Even singularity $X_{e,5}^{\pm\pm}$) *Let the Taylor polynomial of the 4th order of an even function $f(x, y)$ at the point 0 have the form $\pm x^4 + ax^2y^2 \pm y^4$ where $|a| \neq 2$ in the cases $++$ and $--$. Then the germ of f at zero is R_O -equivalent to its Taylor polynomial of the 4th degree.*

A proof to this theorem is analogous to the proof of a similar assertion for arbitrary smooth functions, see the theorem from [1, v. 1, 12.6] about a normal form of semiquasihomogeneous singularities. Here one should use the Tougeron Theorem 3.7 for even functions.

Let us analyze case 2. We will use the Newton diagram (Fig. 1) which corresponds to the Taylor series $\sum a_{p,q}x^py^q$ of f and is defined as follows. This diagram represents the support $\text{supp } f$ consisting of those integer points $(q, p) \in \mathbb{R}^2$ which are exponents of the monomials contributing to the series with nonvanishing coefficients.

The *method of a ruler turning* by Newton consists of the following steps. At first, one draws a line (a «ruler») through the exponent of a marked monomial (which is $\pm x^2y^2$ in our case) such that




Fig. 1: The ruler-method for an even germ f with $(0, 4), (2, 2) \in \text{supp } f \subset \Omega$

this line separates the origin from the unmarked points of the support. Later, one turns the ruler around the marked exponent (counter-clockwise in our case) until it meets the exponent of another monomial present.

Subcase 1. The ruler meets just one point $(2k, 0)$ of the q -axis. In this case, one proves that all other points of the support do not affect the R_O -equivalence class of the germ f . Namely, the germ reduces to the form $\pm x^4 + ax^2y^2 \pm y^{2k}$ with $a \neq 0$ (the even singularity $X_{e,k+3}$), see the corollary from [1, v. 1, 12.7]. Moreover, it follows directly from the proof (the cross-words method) that this reduction can be fulfilled by means of an odd change of variables. Here one should use the Tougeron Theorem 3.7 for even functions.

Subcase 2. The ruler meets two points $(k+1, 1)$ and $(2k, 0)$ which correspond to terms xy^{k+1} and y^{2k} with k even. In this case, let us consider the polynomial which is defined by the monomial $\pm x^4$ and the points falling on the ruler:

$$Ax^4 + Bx^2y^2 + Cxy^{k+1} + Dy^{2k} = Ax^4 + y^2 \left[Bx^2 + Cxy^{k-1} + Dy^{2(k-1)} \right].$$

We will call this polynomial the *mean part* of the germ being investigated. It is easy to observe that, by means of an *odd* change of variables $x' = x + \lambda y^{k-1}$, one can turn the coefficient C into zero. Now the subcase 2 is reduced to the subcase 1.

Case 3 is analyzed similarly to case 2 (the ruler turns clockwise at first and counter-clockwise later). Here, by means of an odd change of variables, the germ reduces to the form $\pm x^{2k} + ax^2y^2 \pm y^{2l}$ with $a \neq 0$ (the even singularity $Y_{e,k,l}$).

Let us return to case 1. It remains to analyze the subcase where the 4-jet of the even germ f reduces to the form $\pm(x^2 + y^2)^2$. Let us associate to the even Taylor series $\pm(x^2 + y^2)^2 + \sum_{p+q \geq 6} a_{p,q}x^p y^q$ of f the series of complex numbers $a_3, a_4, \dots$, which are defined as $a_k = \sum_{p=0}^{2k} i^p a_{p, 2k-p}$. In particular, $a_3 = a_{0,6} - a_{2,4} + a_{4,2} - a_{6,0} + i(a_{1,5} - a_{3,3} + a_{5,1})$. Using the Tougeron Theorem 3.7 for even functions, it is easy to check that, by means of an odd change of variables, the germ reduces to the form $\pm(x^2 + y^2)^2 + ay^6$ with $a = |a_3|$ as soon as $a_3 \neq 0$ (the even singularity $\tilde{Y}_{e,3}$). Moreover, one easily shows that there exists a sequence of polynomials $\tilde{a}_3 = a_3, \tilde{a}_4, \dots$ in $a_{p,q}$ with complex coefficients such that, by means of an odd change of variables, the germ f reduces to the form $\pm(x^2 + y^2)^2 + ay^{2k}$ with $a = |\tilde{a}_k|$ as soon as k is the index of the first nonvanishing element of this sequence (the even singularity $\tilde{Y}_{e,k}$). Here the polynomial $\tilde{a}_k - a_k$ explicitly depends only on $a_{p,q}$, $6 \leq p+q < 2k$. These properties of the sequence $a_3, \tilde{a}_4, \tilde{a}_5, \dots$ can be considered as an analogue of the method of a ruler turning, see above. Let us notice that we actually do not need to consider the case $a_3 = 0$, since the codimension of the corresponding subspace of the space of even singularities having value 0 at the origin equals 7. Here, one considers the subspace consisting of those singularities which satisfy the following condition: the 4-jet of the singularity reduces to the form $\pm(x^2 + y^2)^2$ by means of an odd change of variables and, in the obtained variables, $a_3 = 0$.

Case 4. It is easy to show that, by means of an odd change of variables, the 6-jet of the germ

reduces to the form $x^3y + a_1y^6 + a_2xy^5$. Moreover, the germ itself reduces to the form $x^3y \pm y^6 + axy^5$ if $a_1 \neq 0$ (the even singularity $Z_{e,7}$). Consider the space of even germs in two variables such that the Taylor polynomial of the fourth order reduces to the form x^3y by means of an odd change of variables, and the coefficient at the term y^6 (with respect to these variables) vanishes. It is easy to check that the validity of this condition does not depend on the choice of variables. The codimension of this subspace in the space of all even germs taking value 0 at the origin equals 6. In fact, this subspace is defined by three equations on the 2nd order terms of the Taylor series, two equations on the 4th order terms, and one equation on the coefficient at the term y^6 .

4 Even deformations of even functions

Definition 4.1 An *even deformation* of an even germ $f \in \mathfrak{m}_e$ with a *base* $\Lambda \subset \mathbb{R}^l$ is the germ of a smooth map $F : (\mathbb{R}^\nu \times \Lambda, 0) \rightarrow (\mathbb{R}, 0)$ such that $F(x, 0) \equiv f(x)$ and $F(\cdot, \lambda) \in \mathfrak{m}_e$ for an arbitrary value of the parameter $\lambda \in \Lambda$.

Definition 4.2 An even deformation F' is *R_O -equivalent* to an even deformation F if

$$F'(x, \lambda) \equiv F(g(x, \lambda), \lambda)$$

where $g : (\mathbb{R}^\nu \times \mathbb{R}^l, 0) \rightarrow (\mathbb{R}^\nu, 0)$ is a smooth germ which is odd with respect to x and satisfies the condition $g(x, 0) = x$.

Definition 4.3 A deformation F' is *induced* from F if

$$F'(x, \lambda') \equiv F(x, \varphi(\lambda'))$$

where $\varphi : (\mathbb{R}^l, 0) \rightarrow (\mathbb{R}^l, 0)$ is the germ of a smooth map between bases.

Definition 4.4 An even deformation F of an even germ $f \in \mathfrak{m}_e$ is called *R_O -versal* if any even deformation of f is R_O -equivalent to a deformation which is induced from F . Therefore, an even deformation F of f is R_O -versal if any even deformation F' of f can be represented as

$$F'(x, \lambda') \equiv F(g(x, \lambda'), \varphi(\lambda')), \quad g(x, 0) = x, \quad \varphi(0) = 0, \quad g(-x, \lambda) = -g(x, \lambda). \quad (4.5)$$

With regard to 4.8, the following theorems provide a means for the investigation of singularities of even germs and caustic surfaces.

Theorem 4.6 Any R_O -versal even deformation of a finite-even-multiple even germ f has at least $\mu_e - 1$ parameters, where μ_e is the even multiplicity of f . An even deformation

$$F(x, \lambda) = f(x) + \lambda_1 e_1(x) + \dots + \lambda_{\mu_e-1} e_{\mu_e-1}(x)$$

is R_O -versal if the germs of functions e_k at zero define a basis of the linear space $Q_{\nabla f}^e$.

Theorem 4.7 Any l -parameter R_O -versal even deformation of an even germ f is R_O -equivalent to an even deformation which is induced from any other R_O -versal even deformation with l parameters by means of a diffeomorphism of bases.

Proofs are similar to the proofs of theorems from [1, v. 1, 8.2 and 8.3] and [1, v. 1, 8.5], resp. Actually, Theorems 3.7, 4.6, and 4.7 are partial cases of the corresponding equivariant theorems. The equivariant theorems can be proved in the same way as their non-equivariant analogues, since the groups of germs of equivariant changes of variables in $\mathbb{R}^\nu$ satisfy the conditions of J. Damon [3, 3.2.4 and 3.2.5].

4.8 Stable families of even functions

Let us make an important remark relating the theory of germs deformations and the perturbation theory for parametric families of functions.

Let U be a domain of $\mathbb{R}^n$ which contains 0 and is invariant under the transformation $x \mapsto -x$, and let Λ be a domain of $\mathbb{R}^l$ which contains 0. A smooth function $F: U \times \Lambda \rightarrow \mathbb{R}$ is called a *smooth family of even functions on U with the base Λ* if $F(\cdot, \lambda) \in \mathfrak{m}_e$ for any value of the parameter $\lambda \in \Lambda$, compare 4.1. Any representative $F: (U \times \Lambda, 0) \rightarrow (\mathbb{R}, 0)$ of an even deformation of an even germ $f \in \mathfrak{m}_e$ is obviously a smooth family of even functions.

Remark 4.9 The *R -versality of a deformation* is equivalent to the stronger property *R -stability of a deformation*, see Remark 3 from [1, v. 1, 8.4] (compare the definition from [1, v. 1, 21.1]). Similarly, the *R_O -versality of an even deformation* is equivalent to the stronger property *R_O -stability of an even deformation*: any representative $F: (U \times \Lambda, 0) \rightarrow (\mathbb{R}, 0)$ of an even deformation F admits a neighbourhood E in the space of smooth families of smooth even functions on U with the base Λ (i.e. smooth functions on $U \times \Lambda$ satisfying the conditions $F(-x, \lambda) = F(x, \lambda)$ and $F(0, \lambda) = 0$) such that, for any family $F': U \times \Lambda \rightarrow \mathbb{R}$ from E , there exists a point $0'$ such that the germ F' at $0'$ defines a deformation of a germ f' , R_O -equivalent to the deformation F , where f' is R_O -equivalent to f . Moreover, if the family F' is close enough to F , the point $0'$ can be chosen arbitrarily close to 0 and the R_O -equivalence can be chosen arbitrarily close to the identity.

The versality and the stability of a deformation are apparently equivalent to the stronger property *strict stability*, which is formulated as follows in the case of even deformations: any representative $F: (U \times \Lambda, 0) \rightarrow (\mathbb{R}, 0)$ of an even deformation F admits a neighbourhood E in the space of smooth families of smooth even functions on U with the base Λ and a neighbourhood $U' \times \Lambda'$ of the origin in $U \times \Lambda$ such that U' is invariant under the transformation $(x, \lambda) \mapsto (-x, \lambda)$ and, for any family $F': U \times \Lambda \rightarrow \mathbb{R}$ from E , there exist mappings $g: U' \times \Lambda' \rightarrow U \times \Lambda$ and $\varphi: \Lambda' \rightarrow \Lambda$, close to the identities, such that

$$F(x, \lambda) \equiv F'(g(x, \lambda), \varphi(\lambda)), \quad g(-x, \lambda) = -g(x, \lambda)$$

in the domain $U' \times \Lambda'$. We remark that the conditions $g(x, 0) = x$ and $\varphi(0) = 0$ are not required, compare (4.5).

The property R^+ -stability of an R^+ -versal deformation (resp. R_O -stability of an R_O -versal deformation) implies that a singularity of multiplicity (resp. even multiplicity) μ is non-removable for generic l -parameter families of functions with $l \geq \mu - 1$. Moreover, the caustic surfaces for such families have the same local structure as the caustic surfaces for l -parameter R^+ -versal (resp. R_O -versal) deformations.

Remark 4.10 R^+ -versal deformations of all known types of finite-multiple singularities are R^+ -stratified in the following sense. If $F: (U \times \Lambda) \rightarrow (\mathbb{R}, 0)$ is a representative of an R^+ -versal deformation of a germ $f \in E$ then there exists a neighbourhood $U' \times \Lambda'$ of the origin in $U \times \Lambda$ with the following properties. For any $(x, \lambda) \in U' \times \Lambda' \setminus \{0\}$, the multiplicity of the germ $F(\cdot, \lambda)$ at x is not larger than the multiplicity μ of the germ f at 0. Furthermore, the points $(x, \lambda) \in U' \times \Lambda'$ corresponding to germs which are R^+ -equivalent to the germ f form a smooth submanifold M_1 of codimension $\mu - 1$, while the points $(x, \lambda) \in U' \times \Lambda'$ corresponding to germs of multiplicity μ form a smooth submanifold $M_2 \subset M_1$. Moreover, the restriction of the projection $(x, \lambda) \mapsto \lambda$ to any of the submanifolds M_1 and M_2 is a diffeomorphism onto its image. Here the image of the submanifold M_2 is called a $\mu = \text{const}$ *stratum*, its codimension c is called the *codimension* of the singularity of f , and the integer $m = \mu - c - 1$ is called the *modality* of the singularity of f .

Similarly, R_O -versal deformations of all known types of finite-even-multiple singularities of even germs are R_O -stratified in the following sense. If $F: (U \times \Lambda) \rightarrow (\mathbb{R}, 0)$ is a representative of an R_O -versal deformation of an even germ $f \in \mathfrak{m}_e$ then there exists a neighbourhood $U' \times \Lambda'$ of the origin in $U \times \Lambda$ such that U' is invariant under the transformation $(x, \lambda) \mapsto (-x, \lambda)$ and the following conditions are fulfilled. For any $\lambda \in \Lambda'$, the even multiplicity of the germ $F(\cdot, \lambda)$ at 0 is not larger than the even multiplicity μ_e of the germ f at 0. Moreover, the points $\lambda \in \Lambda'$ corresponding to even germs which are R_O -equivalent to the even germ f form a smooth submanifold of codimension $\mu_e - 1$, while the points $\lambda \in \Lambda'$ corresponding to even germs of even multiplicity μ_e form a smooth submanifold called an *even $\mu_e = \text{const}$ stratum*. Here the codimension c_e of the even stratum is called the *even codimension* of the even singularity of f , and the integer $m_e = \mu_e - c_e - 1$ is called the *even modality* of the even singularity of f .

If a singularity of multiplicity μ admits an R^+ -versal deformation which is R^+ -stratified and strictly R^+ -stable then the class of singularities containing this singularity appears in typical l -parameter families of functions if and only if $l \geq c = \mu - m - 1$.

Similarly, if an even singularity of even multiplicity μ_e admits an R_O -versal deformation which is R_O -stratified and strictly R_O -stable then the even class of singularities containing this singularity appears in typical l -parameter families of even functions if and only if $l \geq c_e = \mu_e - m_e - 1$. **Example.** Consider an even germ $f \in \mathfrak{m}_e$ at the origin with a singularity of type $A_{e,4}$, see Table (2.7). Theorem 4.6 implies that, as an even R_O -versal deformation of f , one can take

$$F(x, \lambda) = \pm x^8 + \lambda_1 x^6 + \lambda_2 x^4 + \lambda_3 x^2.$$

Thus, an even germ of type $A_{e,4}$ becomes non-removable under small perturbations of a family if the dimension l of the parametric space is at least three. Moreover, the surface of caustic values corresponding to singularities of this type has codimension $l - 3$, see 4.10.

Example. Consider an even germ $f \in \mathfrak{m}_e$ at the origin with the even singularity $X_{e,5}^{++}$, see Table (2.7). Theorem 4.6 implies that, as an even R_O -versal deformation of f , one can take

$$F(x, y; \lambda) = x^4 + ax^2y^2 + y^4 + \lambda_1 x^2 + \lambda_2 xy + \lambda_3 y^2 + \lambda_4 x^2y^2.$$

Thus, even germs of class $X_{e,5}^{++}$ become non-removable under small perturbations of an even family if the dimension l of the parametric space is at least four. Observe however that if $\lambda_1 = \lambda_2 = \lambda_3 = 0$ then, for small values of λ_4 , the even germ $F(\cdot; \lambda)$ at zero has a singularity of class $X_{e,5}^{++}$, although it is not equivalent to its initial form $f = F(\cdot; 0)$. Thus, if we want an even singularity of class $X_{e,5}^{++}$ to be preserved under arbitrary small perturbations of an even family, it is enough to require that the dimension l of the parametric space is at least three. Such singularities are called *modal* (in our case, the even singularity of f is unimodal, since it contains one even module a). This term is used for (even) germs whose arbitrary small neighbourhood in the space of (even) germs is covered by a finite number of m -parameter families of (even) singularities. The minimal integer m with this property is called the (even) *modality* of the singularity [1, v. 1, II].

5 Bifurcations of critical points of even functions in typical families

Consider a family $\omega_\lambda(k) : \mathbb{T}^\nu \rightarrow \mathbb{R}$ satisfying (1.1). Values of the parameter λ run through some domain in $\mathbb{R}^l$. We will study *generic* families of functions $\omega_\lambda(k)$, that is such families that all critical points of $\omega_\lambda(\cdot)$ are non-removable for the entire family under small perturbations of the family.

For an arbitrary dimension l of the parametric space, there are only three types of caustic surfaces of codimension one.

1. A caustic hypersurface of type $A_{e,2}^+ \subset A_3$ or $A_{e,2}^- \subset A_3$. It corresponds to a singularity at a

basic critical point. Under a passage of the parameter across this hypersurface, a pair $(k, -k)$ of additional critical points separates from the basic critical point.

2. A caustic hypersurface of type A_2 . It corresponds to a singularity at a twin critical point. Under a passage of the parameter across this hypersurface, two pairs of additional critical points (existing on one side of the hypersurface) merge into one pair and disappear (i.e. are absent on the other side of the hypersurface).

Below, we will give a detailed description of the behaviour (bifurcations) of critical points near the caustics for small (physical) dimensions $l = 1, 2, 3$ of the parametric space. Bifurcations for the codimensions $l = 1, 2$ are described in the works [7, 5].

In Sections 6 and 7, we will show how to determine types of critical points of functions which appear in typical parametric families with a small number $l = 1, 2, 3, 4$ of parameters, and formulate conditions on their deformations to be typical (more precisely, versal).

5.1 Typical one-parameter families

Although we considered even germs taking value 0 at the origin, the values of an even function at the basic critical points freely depend on λ . For convenience, we will assume below that the values of even functions at the basic critical points vanish.

There exists a finite number of caustic values $\lambda_1, \dots, \lambda_p$, for which a *basic* critical point degenerates. Due to Table (2.7) and Statement 6.3, the singularity types at these points are $A_{e,2}$. There exists a finite number of caustic values $\bar{\lambda}_1, \dots, \bar{\lambda}_q$, for which *additional* (twin) critical points degenerate. Due to Table (2.3) and Statement 6.1, the singularity type at these points is A_2 .

Remark 5.2 It is easy to observe that an additional critical point which is a local extremum (i.e. a point of local minimum or local maximum) does not degenerate.

By analyzing (even) versal deformations for degenerate critical points, we understand what happens under a passage of the parameter across a caustic value:

1. $(A_{e,2}^\pm)$ An even versal deformation of a function in a neighbourhood of a degenerate *basic* critical point 0 has the form $F(k, \lambda) = \pm k_1^4 + (\lambda - \lambda_i)k_1^2 \pm k_2^2 \pm \dots \pm k_\nu^2$, see Statement 7.2. Under a passage of the parameter across the caustic value λ_i , a pair $(k, -k)$ of additional critical points separates from the basic critical point.

2. (A_2) An R^+ -versal deformation of a function in a neighbourhood of a degenerate *additional* critical point $\bar{k} = (\bar{k}_1, \dots, \bar{k}_\nu)$ has the form $F(k, \lambda) = (k_1 - \bar{k}_1)^3 + (\lambda - \bar{\lambda}_i)(k_1 - \bar{k}_1) \pm k_2^2 \pm \dots \pm k_\nu^2$, see Statement 7.1. Under a passage of the parameter across the caustic value $\bar{\lambda}_i$, two twin critical points $(\bar{k}', -\bar{k}')$ and $(\bar{k}'', -\bar{k}'')$ which exist for values of λ lying on one side of $\bar{\lambda}_i$ (namely, for $\lambda < \bar{\lambda}_i$) merge into a twin critical point $(\bar{k}, -\bar{k})$ and disappear for λ lying on the other side of $\bar{\lambda}_i$ (namely, for $\lambda > \bar{\lambda}_i$). Bifurcations of level lines of a function for the case $\nu = 2$ are illustrated in Fig. 45 [1, v. 1, 9.6].

5.3 Typical two-parameter families

For $l = 2$, all caustic values form a one-dimensional (not necessarily connected) curve which has one of the following forms in a neighbourhood of any of its points:

1. (A_2) A one-dimensional curve corresponding to a degeneracy of an *additional* critical point with singularity type A_2 . Under a passage of the parameter across this curve, two pairs of critical points merge into one pair $(k_0, -k_0)$ and disappear.

2. $(A_{e,2}^\pm)$ A one-dimensional curve corresponding to a degeneracy of a *basic* critical point with singularity type $A_{e,2} \subset A_3$. Under a passage of the parameter across this curve, a pair $(k, -k)$ of additional critical points separates from the basic critical point.



Fig. 2 **a:** A semicubical cusp point – a typical singularity $A_3^\pm$ of caustic curves
b: Bifurcations of functions at a singularity A_3^+ in typical families of functions




Fig. 3 **a:** A typical singularity $A_{e,3}^\pm$ of caustic curves
b: Bifurcations of functions at a singularity $A_{e,3}^+$ in typical families of even functions

3. ($A_3^\pm$) In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of an *additional* critical point with singularity type A_3 , the set of caustic values consists of two curves Γ_1, Γ_2 of type A_2 (corresponding to a degeneracy of different twin critical points) which adjoin the point $\bar{\lambda}$ (a «point of regression», or a «cusp point») and are tangent to each other at this point with contact order $3/2$ (a «semicubical parabola»), see Fig. 2.

For $\nu = 1$ and $\bar{\lambda} = 0$, an R^+ -versal deformation of a function in a neighbourhood of a degenerate additional critical point $\bar{k}$ of type A_3^+ has the form $F(k, \lambda) = (k - \bar{k})^4 + \lambda_1(k - \bar{k})^2 + \lambda_2(k - \bar{k})$. The caustic curve is defined by the equation $\lambda_2^2 = (\frac{2}{3})^3(-\lambda_1)^3$.

Let us describe the behaviour of critical points for those values of the parameter λ which are close to the point $\bar{\lambda}$: In the domain I , there is one twin critical point $(k_1, -k_1)$. Under a passage across the curve Γ_1 from the domain I to the domain II , two twin critical points $(k'_1, -k'_1)$ and $(k'_2, -k'_2)$ arise. Under a passage across the curve Γ_2 from the domain II to the domain I , the points $(k_1, -k_1)$ and $(k'_1, -k'_1)$ merge and disappear, and the point $(k'_2, -k'_2)$ is renamed $(k_1, -k_1)$.

4. ($A_{e,3}^\pm$) In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degenerate *basic* critical point of type $A_{e,3} \subset A_5$, the set of caustic values consists of two curves Γ_1, Γ_2 of type $A_{e,2} \subset A_3$ (corresponding to a degeneracy of the basic critical point) which adjoin $\bar{\lambda}$ and form a regular curve $\Gamma_1 \cup \bar{\lambda} \cup \Gamma_2$, as well as one curve Γ' of type A_2 (corresponding to a degeneracy of a twin critical point) which also adjoin $\bar{\lambda}$ and is tangent to the curve Γ_2 with the contact order 2, see Fig. 3.

For $\nu = 1$ and $\bar{\lambda} = 0$, an R^+ -versal deformation of a function in a neighbourhood of a degenerate basic critical point 0 of type $A_{e,3}^+$ has the form $F(k, \lambda) = k^6 + \lambda_1 k^4 + \lambda_2 k^2$. The branches $\Gamma_1 \cup \bar{\lambda} \cup \Gamma_2$ and Γ' of the caustic curve are defined by the equations $\lambda_2 = 0$ and resp. $\lambda_2 = \frac{1}{3}\lambda_1^2$, $\lambda_1 < 0$.

Let us describe the behaviour of critical points for those values of the parameter λ which are close to the point $\bar{\lambda}$: In the domain I , there are no twin critical points. Under a passage across the caustic Γ_1 to the domain II , a twin critical point $(k_1, -k_1)$ separates from the basic critical point.

Then, under a passage across the caustic Γ_2 to the domain *III*, one more twin point $(k_2, -k_2)$ separates. Finally, under a passage of λ from the domain *III* to the domain *I* across the caustic Γ' , both twin points merge and disappear.

Of course, except for the singularities listed above, transversal intersections of different branches of caustics are also possible:

5, 6, 7. $(A_2 + A_2, A_2 + A_{e,2}^\pm, A_{e,2}^\pm + A_{e,2}^\pm)$ In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of *two different* (basic or additional) critical points with singularity types A_2 or $A_{e,2}^\pm$, the set of caustic values is the union of two curves Γ_1, Γ_2 (corresponding to a degeneracy of different critical points) which intersect each other transversally at the point $\bar{\lambda}$.

5.4 Typical three-parameter families

All caustic values form a two-dimensional (not necessarily connected) surface which has one of the following forms in a neighbourhood of any of its points:

1. $(A_{e,2}^\pm)$ A surface corresponding to a degeneracy of a *basic* critical point with singularity type $A_{e,2} \subset A_3$. Under a passage of the parameter across this surface, a twin critical point $(k, -k)$ separates from the basic critical point.
2. (A_2) A surface corresponding to a degeneracy of an *additional* critical point with singularity type A_2 . Under a passage of the parameter across this surface, two twin critical points $(k_1, -k_1)$ and $(k_2, -k_2)$ merge into one twin point $(k_0, -k_0)$ and disappear.
3. $(A_3^\pm)$ Two caustic surfaces of type A_2 corresponding to a degeneracy of an *additional* point and adjoining a curve of type A_3 (an «edge of regression», or a «cuspidal edge», see Fig. 55 [1, v. 1, 21.3]). Bifurcations of critical points happen similarly to those in the case of two-parameter even families, see (A_3) .
4. $(A_{e,3}^\pm)$ Three caustic surfaces of types $A_2, A_{e,2}, A_{e,2} \subset A_3$ adjoining a common curve of type $A_{e,3} \subset A_5$ which corresponds to a degeneracy of a *basic* point. Bifurcations of critical points happen similarly to those in the case of two-parameter even families, see $(A_{e,3})$.
5. (A_4) In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of an *additional* critical point with singularity type A_4 , the set of caustic values has the form shown on Fig. 56 [1, v. 1, 21.3] (the «swallow-tail»). Bifurcations of functions are shown in Fig. 28 [2], see also Fig. 55 and 97 [2]).
6. (D_4^+) In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of an *additional* critical point with singularity type D_4^+ , the set of caustic values has the form shown in Fig. 57 [1, v. 1, 21.3] (the «purse»).
7. (D_4^-) In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of an *additional* critical point with singularity type D_4^- , the set of caustic values has the form shown in Fig. 57 [1, v. 1, 21.3] (the «pyramid»).
8. $(A_{e,4}^\pm)$ In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of a *basic* critical point with singularity type $A_{e,4} \subset A_7$, the set of caustic values is formed by the following surfaces (Fig. 4, see also Fig. 62 [1]):
 - a) Two surfaces Γ_1, Γ_2 corresponding to a degeneracy of a *basic* critical point. These surfaces adjoin each other on a curve l_1 which passes through the point $\bar{\lambda}$. Here the degeneracy type of the basic point of this curve equals $A_{e,3} \subset A_5$ (except the point $\bar{\lambda}$). The point $\bar{\lambda}$ splits the curve l_1 into two curves which will be denoted by l_1^+ and l_1^- , resp.
 - b) A surface Γ'_1 corresponding to a degeneracy of an *additional* critical point. The surface Γ'_1 adjoins the surfaces Γ_1, Γ_2 on the curve l_1^- , as well as intersects the surface Γ_1 in a curve l which is tangent to l_1^+ at the point $\bar{\lambda}$. A degeneracy of a basic point of type $A_{e,2} \subset A_3$ and a degeneracy of an additional point of type A_2 correspond to values $\lambda \in l$ (notation $A_{e,2} + A_2$).
 - c) A surface Γ'_2 corresponding to a degeneracy of type A_2 of an *additional* critical point. The




Fig. 4: A typical singularity $A_{e,4}^{\pm}$ of caustic surfaces

surface Γ'_2 adjoins the surfaces Γ_1, Γ_2 on the curve l_1^+ . Moreover the contiguity of Γ'_1 and Γ'_2 to the surfaces Γ_1, Γ_2 takes place on different sides. The surfaces Γ'_1 and Γ'_2 adjoin each other on the curve l'_1 which is tangent to l_1 at the point $\bar{\lambda}$ and corresponds to a degeneracy of type A_3 of an additional critical point.

For $\nu = 1$ and $\bar{\lambda} = 0$, an even versal deformation of a function in a neighbourhood of a degenerate basic critical point 0 of type $A_{e,4}^+$ has the form $F(k, \lambda) = k^8 + \lambda_1 k^6 + \lambda_2 k^4 + \lambda_3 k^2$. The branches $\Gamma_1 \cup l_1 \cup \Gamma_2$ and Γ'_1, Γ'_2 of the caustic surface are respectively defined by the equations

$$\begin{aligned} \lambda_3 &= 0, \\ 8\lambda_3 + \lambda_1(-4\lambda_2 + \lambda_1^2) &= (-\frac{8}{3}\lambda_2 + \lambda_1^2)^{3/2} \quad \text{for } \lambda_2 < 0 \text{ or } 0 \leq \lambda_2 \leq \frac{3}{8}\lambda_1^2, \lambda_1 < 0, \\ 8\lambda_3 + \lambda_1(-4\lambda_2 + \lambda_1^2) &= -(-\frac{8}{3}\lambda_2 + \lambda_1^2)^{3/2} \quad \text{for } 0 < \lambda_2 \leq \frac{3}{8}\lambda_1^2, \lambda_1 < 0. \end{aligned}$$

Bifurcations of even functions in $\nu = 1$ variable in a neighbourhood of a caustic value corresponding to a singularity of type $A_{e,4}^+$ are shown in Fig. 5 (compare Fig. 91 [2]). This figure does not show the whole caustic, but only its intersection with a cylinder ($\lambda_1 = \varepsilon \sin \varphi$, $\lambda_2 = \varepsilon \cos \varphi$, $0 \leq \varphi \leq 2\pi$ for a small positive constant ε), which intersects the plane $\Gamma_1 \cup l_1 \cup \Gamma_2$ transversally. Let us describe the behaviour of critical points for parameter values λ close to $\bar{\lambda}$:

In the domain I , there are no twin critical points. Under a passage across the caustic Γ_1 to the domain II , a twin point $(k_1, -k_1)$ separates from the basic critical point.

Under a passage across the caustic Γ'_1 from the domain II to the domain III , two twin critical points $(k_2, -k_2)$ and $(k_3, -k_3)$ arise. Thus, for the domain III , there are 3 twin critical points in a neighbourhood of the basic critical point.

Under a passage across the caustic Γ'_2 from the domain III to the domain II , the twin critical points $(k_1, -k_1)$ and $(k_2, -k_2)$ merge and disappear.

Under a passage across the caustic Γ_2 from the domain II to the domain IV , the twin critical point $(k_2, -k_2)$ separates from the basic critical point.

Under a passage across the caustic Γ'_1 from the domain IV to the domain I , the twin critical points $(k_2, -k_2)$ and $(k_3, -k_3)$ merge and disappear.

9, 10. ($X_{e,5}^{++}$, $X_{e,5}^{--}$, and $X_{e,5}^{+-}$) As a “model” (more precisely, a one-parameter family of “models”) of a typical 3-parameter family of functions in $\nu = 2$ variables, we consider

$$\pm x^4 + ax^2y^2 \pm y^4 + \lambda_1x^2 + \lambda_2xy + \lambda_3y^2.$$

Here $\lambda_1, \lambda_2, \lambda_3$ are parameters of the family and a is a parameter of the model ($a \neq \pm 2$ in the cases $++$ and $--$). In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of a *basic* critical point with singularity class $X_{e,5} \subset X_9$, the set of caustic values is a surface which is the union of two conic surfaces (more precisely, “surfaces close to conic ones”) Γ and Γ' with vertex at





Fig. 5: Bifurcations of functions at a singularity $A_{e,4}^+$ in typical families of even functions

the point $\bar{\lambda}$. Moreover these surfaces can intersect each other on rays coming from $\bar{\lambda}$. The surface Γ (except the point $\bar{\lambda}$) corresponds to degeneracies of types $A_{e,2}$ and $A_{e,3}$ of a *basic* critical point, while the surface Γ' (except the point $\bar{\lambda}$) corresponds to degeneracies of types A_2 and A_3 of an *additional* critical point. (Actually, for our model, as for any generic 1-parameter family of “models” with a parameter a , degeneracies of types A_4 and $D_4^\pm$ as well as non-transversal intersections of different branches of the caustic can happen in general for some exceptional values of a . However the number of such exceptional values of a is finite, thus the corresponding models are not generic and, hence, one does not need to consider them.) More precisely, the surfaces Γ and Γ' have the following form:

a) The surface Γ is a circular cone having vertex at the point $\bar{\lambda}$ (in our model it is the cone $\lambda_2^2 = 4\lambda_1\lambda_3$). Finitely many elements $l_1, l_2, \dots$ of the cone Γ depending on a are chosen: In the case $++$, there are no chosen elements of the cone for $a > -2$, while there are four chosen elements l_1, l_2, l_3, l_4 for $a < -2$ (for our model, they are obtained by intersecting the cone with two planes $\frac{\lambda_3}{\lambda_1} = \frac{-a+\sqrt{a^2-4}}{2}$ and $\frac{\lambda_3}{\lambda_1} = \frac{-a-\sqrt{a^2-4}}{2}$). Similarly, in the case $--$, there are no chosen elements of the cone for $a < 2$, while there are four chosen elements l_1, l_2, l_3, l_4 for $a > 2$ (for our model, they are obtained by intersecting the cone with two planes $\frac{\lambda_1}{\lambda_3} = \frac{a+\sqrt{a^2-4}}{2}$ and $\frac{\lambda_1}{\lambda_3} = \frac{a-\sqrt{a^2-4}}{2}$). In the case $+-$ two elements l_1 and l_2 of the cone are chosen (for our model, they are obtained by intersecting the cone by the plane $\frac{\lambda_1}{\lambda_3} = \frac{-a+\sqrt{a^2+4}}{2}$).

All points of the cone Γ except the point $\bar{\lambda}$ and the points lying on the chosen cone elements $l_1, l_2, \dots$ correspond to a degeneracy of type $A_{e,2}$ of a *basic* critical point. All points of the chosen cone elements apart from $\bar{\lambda}$ correspond to a degeneracy of type $A_{e,3}$ of a *basic* critical point.

Denote by Γ^+ and Γ^- two half-cones of the cone Γ , i.e. two connected components of $\Gamma \setminus \bar{\lambda}$ (for our model, they are the parts of Γ lying in the octants $\lambda_1 > 0, \lambda_3 > 0$, and resp. $\lambda_1 < 0, \lambda_3 < 0$).



Furthermore, in the case $+-$ with $a = 0$, the surface Γ' is the closure of the surface defined by the equation

$$(\varrho - 1)^3 = \frac{27}{16}(2 + \sigma)\varrho \quad \begin{array}{l} \text{for } \varrho < 0, \lambda_1 < 0, \\ \text{and for } \varrho \geq 4, \lambda_1(\lambda_1^2 - \lambda_3^2) < 0. \end{array}$$

Here one denotes $\varrho = \frac{\lambda_2^2}{\lambda_1\lambda_3}$, $\sigma = \frac{\lambda_1^2}{\lambda_3^2} + \frac{\lambda_3^2}{\lambda_1^2}$. Actually, in the case $+-$, the conic surface Γ' is obtained from the indicated surface by adding four rays, where two of these rays lie on the plane $\lambda_1 = 0$, and the other two rays lie on the plane $\lambda_3 = 0$.

In the cases $++$ and $+-$ with $0 < a \ll 1$, bifurcations of level lines of functions in $\nu = 2$ variables are shown on Fig. 7 and 8, respectively. These figures do not show the whole caustic, but only its intersection with the cylinder $\lambda_1 = \varepsilon \cos \varphi$, $\lambda_3 = \varepsilon \sin \varphi$, $-\frac{\pi}{4} \leq \varphi \leq \frac{7\pi}{4}$ for a small positive constant ε . In the both cases, the bifurcation diagram is invariant under the transformation $(\lambda_2, \varphi, x, y) \mapsto (-\lambda_2, \varphi, -x, y)$. One can see from these figures that, if the parameter λ goes around a loop which envelops the cone Γ , the following transformation happens. In the case $++$, the twin critical point $(k, -k)$ (corresponding to a pair of minima of an even function) transforms into $(-k, k)$. In the case $+-$, each of two separatrices of the saddle point 0 turns by π .

Remark 5.5 The conic surface Γ' is symmetric. Namely, it is invariant under reflections with respect to one or two planes which leave the cone Γ invariant. In our model, it is the plane $\lambda_2 = 0$ in the case $+-$ and two planes $\lambda_2 = 0$ and $\lambda_1 = \lambda_3$ in each of the cases $++$ and $--$. Besides, in the case $+-$, the caustic surfaces corresponding to opposite values of the parameter a are obtained from each other by reflection with respect to the plane $\lambda_1 = -\lambda_3$. Furthermore, the caustic surfaces in the case $++$ and the case $--$, corresponding to opposite values of the parameter a , are obtained from each other by the reflection with respect to this plane.

Remark 5.6 If the parameter a continuously varies in the domain of admissible values ($a \neq \pm 2$ in the cases $++$ and $--$), the conic surface Γ' is deformed. In particular, locations of the cuspidal edges l'_1, l'_2, l'_3, l'_4 (in the cases $++$ and $--$) and resp. l'_1, l'_2 (in the case $+-$) are continuously deformed with respect to a . Moreover, each of these cuspidal edges is preserved for all admissible values of a by just deforming continuously on a , with the edge remaining in the corresponding plane of symmetry:

In the cases $++$ and $--$, the cuspidal edges l'_1 and l'_3 lie in the plane $\lambda_2 = 0$, while the cuspidal edges l'_2 and l'_4 lie in the plane $\lambda_1 = \lambda_3$. More precisely: in the case $++$, the cuspidal edges l'_1 and l'_3 correspond to the twin critical points $\left((\sqrt{-\frac{\lambda_1}{2}}, 0), (-\sqrt{-\frac{\lambda_1}{2}}, 0) \right)$ and resp.

$\left((0, \sqrt{-\frac{\lambda_3}{2}}), (0, -\sqrt{-\frac{\lambda_3}{2}}) \right)$ of type A_3 . They are defined by $\lambda_3 = \frac{a}{2}\lambda_1$, $\lambda_1 < \lambda_2 = 0$ and resp. $\lambda_1 = \frac{a}{2}\lambda_3$, $\lambda_3 < \lambda_2 = 0$. Furthermore, the cuspidal edges l'_2 and l'_4 correspond to the twin critical points $\left((\sqrt{-\frac{\lambda_1}{2}}, \sqrt{-\frac{\lambda_1}{2}}), (-\sqrt{-\frac{\lambda_1}{2}}, -\sqrt{-\frac{\lambda_1}{2}}) \right)$ and resp. $\left((\sqrt{-\frac{\lambda_1}{2}}, -\sqrt{-\frac{\lambda_1}{2}}), (-\sqrt{-\frac{\lambda_1}{2}}, \sqrt{-\frac{\lambda_1}{2}}) \right)$ of type A_3 . They are defined by $\lambda_2 = \frac{a-2}{2}\lambda_1$, $\lambda_1 = \lambda_3 < 0$ and resp. $\lambda_2 = \frac{2-a}{2}\lambda_1$, $\lambda_1 = \lambda_3 < 0$.

In the case $+-$, the cuspidal edges l'_1 and l'_2 lie in the plane $\lambda_2 = 0$ and correspond to the twin critical points $\left((\sqrt{-\frac{\lambda_1}{2}}, 0), (-\sqrt{-\frac{\lambda_1}{2}}, 0) \right)$ and resp. $\left((0, \sqrt{\frac{\lambda_3}{2}}), (0, -\sqrt{\frac{\lambda_3}{2}}) \right)$ of type A_3 . They are defined by $\lambda_3 = \frac{a}{2}\lambda_1$, $\lambda_1 < \lambda_2 = 0$ and resp. $\lambda_1 = -\frac{a}{2}\lambda_3$, $\lambda_3 > \lambda_2 = 0$.

>From the indicated explicit presentation of the cuspidal edges l'_1, l'_2 (in the case $+-$) and l'_1, l'_2, l'_3, l'_4 (in the cases $++$ and $--$), we find their location with respect to the cone Γ for any a : In the case $++$ (the case $--$ is analogous due to 5.5), all of the cuspidal edges l'_1, l'_2, l'_3, l'_4 remain inside the domain Ω^- for $0 < a < 2$ and for $2 < a < 6$, while they are outside the domains Ω^+ and Ω^- for $a < -2$ (moreover, all these cuspidal edges “tend” to the axis of the cone Γ as $a \rightarrow 2$).




Fig. 7: Bifurcations of functions at a singularity $X_{e,5}^{++}$ with module $a \in \mathbb{R} \setminus \{2, -2\}$ in typical families of even functions, $0 < a \ll 1$




Fig. 8: Bifurcations of functions at a singularity $X_{e,5}^{+-}$ with module $a \in \mathbb{R}$ in typical families of even functions, $0 < a \ll 1$

In the case $++$ with $-2 < a < 0$, the cuspidal edges l'_2 and l'_4 lie inside the domain Ω^- , while the cuspidal edges l'_1 and l'_3 lie outside the domains Ω^+ and Ω^- .

In the case $++$ with $a > 6$, the cuspidal edges l'_1 and l'_3 lie inside the domain Ω^- , while the cuspidal edges l'_2 and l'_4 lie outside the domains Ω^+ and Ω^- .

In the case $+ -$ with $a > 0$, the cuspidal edge l'_1 lies inside the domain Ω^- , while the cuspidal edge l'_2 lies outside the domains Ω^+ and Ω^- .

In the case $+ -$ with $a < 0$, the cuspidal edge l'_2 lies inside the domain Ω^+ , while the cuspidal edge l'_1 lies outside the domains Ω^+ and Ω^- .

Unfortunately, the authors do not know whether the form of the conic surface Γ' (in particular, the number of its cuspidal edges) and its location with respect to the cone Γ change under a continuous change of the value of a . In particular, we do not know whether *bifurcations of cuspidal edges* happen for some exceptional values of a lying outside the interval $0 < |a| \ll 1$ considered above. Possible bifurcations of cuspidal edges should be similar to bifurcations of cusps occurring in generic families of sections by parallel planes of a swallow-tail (a rise/destruction of a pair of cuspidal edges), a purse (a bifurcation of one cuspidal edge), and a pyramid (a bifurcation of three cuspidal edges).

Of course, except for the singularities listed above, transversal intersections of different branches of caustics are also possible:

11, 12. $(A_2 + A_3^\pm, A_{e,2}^\pm + A_3^\pm)$ In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of an *additional* critical point of degeneracy type $A_3^\pm$ and another *basic or additional* critical point of degeneracy type A_2 or resp. $A_{e,2}^\pm$, the set of caustic values is the union of a surface from case 3 ($A_3^\pm$) and a plane (from case 2 or resp. 1) which intersects the exceptional curve of type $A_3^\pm$ of the first surface transversally.

13, 14. $(A_2 + A_{e,3}^\pm, A_{e,2}^\pm + A_{e,3}^\pm)$ In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of a *basic* critical point of degeneracy type $A_{e,3}^\pm$ and another *basic or additional* critical point of degeneracy type A_2 or resp. $A_{e,2}^\pm$, the set of caustic values is the union of a surface from the case 4 ($A_{e,3}^\pm$) and a plane (from the case 2 or resp. 1) which intersects the exceptional curve of type $A_{e,3}^\pm$ of the first surface transversally.

15, 16, 17, 18. $(A_2 + A_2 + A_2, A_2 + A_2 + A_{e,2}^\pm, A_2 + A_{e,2}^\pm + A_{e,2}^\pm, A_{e,2}^\pm + A_{e,2}^\pm + A_{e,2}^\pm)$ In a neighbourhood of a caustic value $\bar{\lambda}$ corresponding to a degeneracy of three different *basic or additional* critical points of degeneracy types A_2 or $A_{e,2}^\pm$, the set of caustic values is the union of three planes which intersect each other transversally at the point $\bar{\lambda}$.

6 A determinant of singularity types

In practice, the following questions often arise: whether the germ of a given function f has a certain singularity type, and which normal form the function has near the singular point. For singularities of codimension 0 (i.e. Morse functions and even Morse functions), answers are given by the Morse lemma for arbitrary functions and even functions, see Theorem 3.5.

Below, we answer these questions for the singularities which appear in typical l -parameter families with $l = 1, 2, 3, 4$ parameters for arbitrary functions and even functions, see Statements 6.1, 6.7, 6.8, and resp. 6.3, 6.10, 6.12.

Statement 6.1 (A singularity A_μ) *Let 0 be a critical point of a smooth function $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$. Suppose that the kernel of the second differential of f at the origin coincides with the coordinate axis Ok_1 (in particular, 0 is a critical point of co-rank 1). Then there exists a sequence of real numbers a_μ , $\mu = 2, 3, \dots$, depending on partial derivatives of f at the origin, and possessing the following properties:*

1. $a_\mu = \frac{\partial^{\mu+1} f(0)}{\partial k_1^{\mu+1}} + \dots$, where the additional terms form a polynomial in the values of the partial derivatives $\frac{\partial^{i_1+\dots+i_\nu} f(0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu}}$, $3 \leq i_1 + \dots + i_\nu \leq \mu$ (except the derivatives with respect to the variable k_1 only) and the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 2, \dots, \nu$. This polynomial has integer coefficients and a vanishing free term. The numbers a_μ are given in Table (6.5) for $\mu = 2, 3, 4, 5$.

2. The germ of f at the origin has a singularity of type A_μ if and only if

$$a_2 = \dots = a_{\mu-1} = 0, \quad a_\mu \neq 0. \quad (6.2)$$

Moreover, if (6.2) is fulfilled then f reduces to the form $f = a + a_\mu \tilde{k}_1^{\mu+1} + Q(\tilde{k}_2, \dots, \tilde{k}_\nu)$ in some neighbourhood of the origin by means of a regular change of variables $k \rightarrow \tilde{k}$ leaving the origin fixed. Here a is a constant, Q is a nondegenerate quadratic form in $\nu - 1$ variables.

In particular, $f \in A_\mu^+$ if $a_\mu > 0$; $f \in A_\mu^-$ if $a_\mu < 0$.

Statement 6.3 (An even singularity $A_{e,\mu}$) Let $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$ be a smooth even function defined in a neighbourhood of the point 0 in $\mathbb{R}^\nu$. Suppose that the kernel of the second differential of f at the origin coincides with the coordinate axis Ok_1 (in particular, 0 is a critical point of co-rank 1). Then there exists a sequence of real numbers $a_{e,\mu}$, $\mu = 2, 3, \dots$, depending on partial derivatives of f at the origin, and possessing the following properties:

1. $a_{e,\mu} = a_{2\mu-1} = \frac{\partial^{2\mu} f(0)}{\partial k_1^{2\mu}} + \dots$ where $a_{2\mu-1}$ is assigned according to Statement 6.1 to the even function f , which is considered as a usual function. The numbers $a_{e,\mu}$ are given in Table (6.5) for $\mu = 2, 3, 4, 5$.

2. The even germ of f at the origin has an even singularity of type $A_{e,\mu}$ if and only if

$$a_{e,2} = \dots = a_{e,\mu-1} = 0, \quad a_{e,\mu} \neq 0. \quad (6.4)$$

Moreover, if (6.4) is fulfilled then f reduces to the form $f = a + a_{e,\mu} \tilde{k}_1^{2\mu} + Q(\tilde{k}_2, \dots, \tilde{k}_\nu)$ in some neighbourhood of the origin by means of a regular odd change of variables $k \rightarrow \tilde{k}$. Here a and Q are the same as in Statement 6.1.

In particular, $f \in A_{e,\mu}^+$ if $a_{e,\mu} > 0$; $f \in A_{e,\mu}^-$ if $a_{e,\mu} < 0$.

μ	a_μ	$a_{e,\mu}$
2	f_{y^3}	f_{y^4}
3	$f_{y^4} - 3f_{z^2}^{-1}[f_{y^2z}]^2$	$f_{y^6} - 10f_{z^2}^{-1}[f_{y^3z}]^2$
4	$f_{y^5} - f_{y^2z}f_{z^2}^{-1}(10f_{y^3z} - 15f_{yz^2}f_{z^2}^{-1}f_{y^2z})$	$f_{y^8} - 56f_{y^3z}f_{z^2}^{-1}(f_{y^5z} - 5f_{y^2z^2}f_{z^2}^{-1}f_{y^3z})$
5	$f_{y^6} - 15f_{y^2z}f_{z^2}^{-1}(f_{y^4z} - 4f_{yz^2}f_{z^2}^{-1}f_{y^3z})$ $+ 6[f_{yz^2}f_{z^2}^{-1}]^2f_{y^2z} - 10f_{z^2}^{-1}[f_{y^3z}]^2$ $+ 45f_{y^2z^2}[f_{z^2}^{-1}f_{y^2z}]^2 - 15f_{z^3}[f_{z^2}^{-1}f_{y^2z}]^3$	$f_{y^{10}} - 120f_{y^3z}f_{z^2}^{-1}(f_{y^7z} - 21f_{y^2z^2}f_{z^2}^{-1}f_{y^5z})$ $+ 105[f_{y^2z^2}f_{z^2}^{-1}]^2f_{y^3z} - 126f_{z^2}^{-1}[f_{y^5z}]^2$ $+ 2100f_{y^4z^2}[f_{z^2}^{-1}f_{y^3z}]^2 - 2620f_{y^3z^3}[f_{z^2}^{-1}f_{y^3z}]^3$

(6.5)

Comment to Table (6.5). In this table, the variables are denoted by $y = k_1$ and $z = (k_2, \dots, k_\nu)$, moreover $f_{y^a z^b} := \frac{\partial^{a+b} f(0)}{\partial y^a \partial z^b}$. The formulae in the table are written for $\nu = 2$, but they are easily transformed to the corresponding formulae for any $\nu \geq 1$. For example:

$$a_\mu = \frac{\partial^{\mu+1} f(0)}{\partial k_1^{\mu+1}} \quad \text{for } \nu = 1, \quad a_3 = \frac{\partial^4 f(0)}{\partial k_1^4} - 3 \sum_{i=2}^{\nu} \sum_{j=2}^{\nu} \frac{\partial^3 f(0)}{\partial k_1^2 \partial k_i} A_{ij} \frac{\partial^3 f(0)}{\partial k_1^2 \partial k_j} \quad \text{for } \nu \geq 2 \quad (6.6)$$

where A_{ij} , $i, j = 2, \dots, \nu$ denote the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 2, \dots, \nu$.

We will prove Statement 6.1 using the technique of the proof of the lemma about classifying finite-multiple singularities of co-rank 1 [1, v. 1, 9.6]. Due to the parametric Morse lemma for arbitrary smooth functions (analogous to Theorem 3.6), in a small neighbourhood of the origin, there exists a smooth change of variables $(k_1, k_2, \dots, k_\nu) \rightarrow (k_1, \tilde{k}_2, \dots, \tilde{k}_\nu)$ leaving the origin fixed and reducing f to the form $f = a + \psi(k_1) + Q(\tilde{k}_2, \dots, \tilde{k}_\nu)$, where a is a constant, Q is a nondegenerate quadratic form, and $\psi(k_1) = O(|k_1|^3)$ as $k_1 \rightarrow 0$. It is obvious that the germ at the origin of a function ψ in one variable has a singularity of type A_μ if and only if the order of the function ψ at the origin is $\mu + 1$, i.e. $\psi'(0) = \psi''(0) = \dots = \psi^{(\mu)}(0) = 0$ and $\psi^{(\mu+1)}(0) \neq 0$.

Therefore, it remains to express the numbers $a_\mu := \psi^{(\mu+1)}(0)$ in terms of the values of the partial derivatives of f at the origin. Observe that $\psi(k_1) = f(k_1, k_2(k_1), \dots, k_\nu(k_1))$ where the functions $k_i = k_i(k_1)$, $i = 2, \dots, \nu$ define the coordinate axis Ok_1 of the coordinate system $k_1, \tilde{k}_2, \dots, \tilde{k}_\nu$ and, hence, are implicitly defined by the system of equations $\frac{\partial f}{\partial k_i}(k_1, k_2(k_1), \dots, k_\nu(k_1)) = 0$, $i = 2, \dots, \nu$. One sets the values at the origin, of the derivatives of order $\leq \mu - 1$ of the left-hand sides of these equations, to zero. This will give expressions for $k'_i(0), \dots, k_i^{(\mu-1)}(0)$ in terms of the partial derivatives of order $\leq \mu$ of f at the origin. By substituting these expressions into the derivative of order $\mu + 1$ at the origin of the composite function ψ , we obtain $a_\mu = \psi^{(\mu+1)}(0) = \frac{\partial^{\mu+1} f(0)}{\partial k_1^{\mu+1}} + \dots$ where «...» is expressed by means of the partial derivatives of order $\leq \mu$ of f at the origin.

Statement 6.3 is similarly proved based on Theorem 3.6.

Statement 6.7 (A singularity D_4) *Let 0 be a critical point of a smooth function $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$. Suppose that the kernel of the second differential of f at the origin coincides with the coordinate plane Ok_1k_2 (in particular, 0 is a critical point of co-rank 2). The germ of f at the origin has a singularity of type D_4 if and only if $\Delta_3 \neq 0$, where Δ_3 is the discriminant*

$$\Delta_3 = 4(B^3D + AC^3) + 27A^2D^2 - B^2C^2 - 18ABCD,$$

$$A = \frac{1}{3!} \frac{\partial^3 f(0)}{\partial k_1^3}, \quad B = \frac{1}{2!} \frac{\partial^3 f(0)}{\partial k_1^2 \partial k_2}, \quad C = \frac{1}{2!} \frac{\partial^3 f(0)}{\partial k_1 \partial k_2^2}, \quad D = \frac{1}{3!} \frac{\partial^3 f(0)}{\partial k_2^3}.$$

Moreover, if $\Delta_3 \neq 0$ then f reduces to the form $f = d + \tilde{k}_1^2 \tilde{k}_2 + \Delta_3 \tilde{k}_2^3 + Q(\tilde{k}_3, \dots, \tilde{k}_\nu)$ in some neighbourhood of the origin by means of a regular change of variables $k \rightarrow \tilde{k}$ leaving the point 0 fixed. Here d is a constant, Q is a nondegenerate quadratic form in $\nu - 2$ variables.

In particular, $f \in D_4^+$ if $\Delta_3 > 0$; $f \in D_4^-$ if $\Delta_3 < 0$.

Statement 6.8 (A singularity D_μ) *Let 0 be a critical point of a smooth function $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$. Suppose that the kernel of the second differential of f at the origin coincides with the coordinate plane Ok_1k_2 (in particular, 0 is a critical point of co-rank 2). Suppose that $\frac{\partial^3 f(0)}{\partial k_1^3} = \frac{\partial^3 f(0)}{\partial k_1 k_2^2} = 0$ and $\frac{\partial^3 f(0)}{\partial k_1^2 \partial k_2} = 1$ (this can be achieved by means of a suitable linear change of the variables k_1, k_2 provided that the cubic part of the Taylor series at the origin of the function $f(k_1, k_2, 0, \dots, 0)$ does not vanish and is not a perfect cube). Then there exists a sequence of real numbers d_μ , $\mu = 4, 5, \dots$, depending on partial derivatives of f at the origin, and possessing the following properties:*

1. $d_\mu = \frac{\partial^{\mu-1} f(0)}{\partial k_2^{\mu-1}} + \dots$, where the additional terms form a polynomial in the values of the partial derivatives $\frac{\partial^{i_1+\dots+i_\nu} f(0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu}}$, $3 \leq i_1 + \dots + i_\nu \leq \mu - 2$ (except the derivatives with respect to the variable k_2 only) and the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 3, \dots, \nu$. This polynomial has rational coefficients and a vanishing free term. The numbers d_μ are given in Table (6.14) for $\mu = 4, 5, 6, 7, 8$.

2. For $\mu \geq 4$, the germ of f at the origin has a singularity of type $D_\mu^\pm$ if and only if

$$d_4 = \dots = d_{\mu-1} = 0, \quad d_\mu \neq 0. \quad (6.9)$$

Moreover, if (6.9) holds then f reduces to the form $f = d + \tilde{k}_1^2 \tilde{k}_2 + d_\mu \tilde{k}_2^{\mu-1} + Q(\tilde{k}_3, \dots, \tilde{k}_\nu)$ in some neighbourhood of the origin by means of a regular change of variables $k \rightarrow \tilde{k}$ leaving the origin fixed. Here d and Q are similar to those in Statement 6.7.

In particular, $f \in D_\mu^+$ if $d_\mu > 0$; $f \in D_\mu^-$ if $d_\mu < 0$.

Statement 6.10 (An even singularity $X_{e,5}$) Suppose that the kernel of the second differential at the origin of a smooth even function $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$ coincides with the coordinate plane Ok_1k_2 (in particular, 0 is a critical point of co-rank 2). Then the even germ of f at the origin has an even singularity of class $X_{e,5}^{\pm\pm}$ if and only if $\Delta_4 \neq 0$, where Δ_4 is the discriminant

$$\begin{aligned} \Delta_4 = & -4(AC^3D^2 + B^3D^3 + B^2C^3E) - 27(A^2D^4 + B^4E^2) + B^2C^2D^2 + 18(ABCD^3 + B^3CDE) \\ & + 144(A^2CD^2E + AB^2CE^2) - 6AB^2D^2E - 80ABC^2DE - 192A^2BDE^2 + 16AE(C^2 - 4AE)^2, \\ A = & \frac{1}{4!} \frac{\partial^4 f(0)}{\partial k_1^4}, \quad B = \frac{1}{3!} \frac{\partial^4 f(0)}{\partial k_1^3 \partial k_2}, \quad C = \frac{1}{4} \frac{\partial^3 f(0)}{\partial k_1^2 \partial k_2^2}, \quad D = \frac{1}{3!} \frac{\partial^3 f(0)}{\partial k_1 \partial k_2^3}, \quad E = \frac{1}{4!} \frac{\partial^4 f(0)}{\partial k_2^4}. \end{aligned}$$

Moreover, $f \in X_{e,5}^{++} \cup X_{e,5}^{--}$ if $\Delta_4 > 0$; $f \in X_{e,5}^{+-}$ if $\Delta_4 < 0$.

Remark 6.11 Suppose that, under hypothesis of Statement 6.10, $\Delta_4 \neq 0$. This is equivalent to the fact that the polynomial $Ax^4 + Bx^3 + Cx^2 + Dx + E$ does not have multiple roots [11]. Therefore, by means of a linear change of variables, the corresponding homogeneous polynomial of the fourth degree in two variables reduces to the form $Ax^4 + Cx^2y^2 + Ey^4$. In these variables, we have $B = D = 0$ and $\Delta_4 = 16AE(C^2 - 4AE)^2$. Consequently, $A \neq 0$, $E \neq 0$, and $C^2 \neq 4AE$. Therefore one can assume that $B = D = 0$, $|A| = |E| = 1$, and that $|C| \neq 2$ as soon as $AE = 1$.

Statement 6.12 (An even singularity $X_{e,\mu}$) Suppose that the kernel of the second differential at the origin of a smooth even function $f : \mathbb{R}^\nu \rightarrow \mathbb{R}$ coincides with the coordinate plane Ok_1k_2 (in particular, 0 is a critical point of co-rank 2). Suppose that

$$\frac{\partial^4 f(0)}{\partial k_1^3 \partial k_2} = \frac{\partial^4 f(0)}{\partial k_1 \partial k_2^3} = 0, \quad \frac{\partial^4 f(0)}{\partial k_1^4} = 24\varepsilon, \quad \frac{\partial^4 f(0)}{\partial k_1^2 \partial k_2^2} = 4\eta, \quad \frac{\partial^4 f(0)}{\partial k_2^4} \neq 6\varepsilon \quad \text{where } \varepsilon, \eta = \pm 1$$

(this can be achieved by means of a suitable linear change of the variables k_1, k_2 provided that the homogeneous polynomial of the 4-th degree $Ak_1^4 + Bk_1^3k_2 + Ck_1^2k_2^2 + Dk_1k_2^3 + Ek_2^4$ of the Taylor series at the origin of the function $f(k_1, k_2, 0, \dots, 0)$ does not vanish, is not a perfect square, and reduces either to the form 1 from 3.12 with $a \neq 0$, or to the form 2 from 3.12). Then there exists a sequence of real numbers $x_{e,\mu} = x_{e,\mu}^{\varepsilon,\eta}$, $\mu = 5, 6, \dots$ depending on partial derivatives of f at the origin and possessing the following properties:

1. $x_{e,\mu}^{\varepsilon,\eta} = \frac{\partial^{2\mu-6} f(0)}{\partial k_2^{2\mu-6}} + \dots$, where the additional terms form a polynomial in the values of the partial derivatives $\frac{\partial^{i_1+\dots+i_\nu} f(0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu}}$, $4 \leq i_1 + \dots + i_\nu \leq 2\mu - 8$ (except the derivatives with respect to the variable k_2 only) and the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 3, \dots, \nu$. This polynomial has rational coefficients and a vanishing free term. The numbers $x_{e,\mu} = x_{e,\mu}^{\varepsilon,\eta}$ are given in Table (6.14) for $\mu = 5, 6, 7, 8$.

2. For $\mu \geq 5$, the even germ of f at the origin has a singularity of class $X_{e,\mu}^{\pm\pm}$ if and only if

$$x_{e,5}^{\varepsilon,\eta} = \dots = x_{e,\mu-1}^{\varepsilon,\eta} = 0, \quad x_{e,\mu}^{\varepsilon,\eta} \neq 0. \quad (6.13)$$

Moreover, if (6.13) holds then f reduces to the form $f = d + \varepsilon \tilde{k}_1^4 + \eta \tilde{k}_1^2 \tilde{k}_2^2 + \frac{x_{e,\mu}^{\varepsilon,\eta}}{(2\mu-6)!} \tilde{k}_2^{2\mu-6} + Q(\tilde{k}_3, \dots, \tilde{k}_\nu)$ in some neighbourhood of the origin by means of a regular odd change of variables $k \rightarrow \tilde{k}$. Here d and Q are similar to those in Statement 6.7.

In particular, $\frac{x_{e,\mu}^{\varepsilon,\eta}}{(2\mu-6)!}$ is the even module of the singularity $X_{e,\mu}^{\varepsilon,\eta}$ if $\mu \geq 6$.

μ	d_μ	$x_{e,\mu} = x_{e,\mu}^{\varepsilon,\eta}$
4	$a_{0,3} = f_{y^3}$	—
5	$a_{0,4} = f_{y^4} - 3f_{z^2}^{-1}[f_{y^2z}]^2$	$a_{0,4} = f_{y^4}$
6	$a_{0,5} = \frac{5}{3}a_{1,3}^2$	$a_{0,6} = f_{y^6} - 10f_{z^2}^{-1}[f_{y^3z}]^2$
7	$a_{0,6} = 5a_{1,3}a_{1,4} + 5a_{1,3}^2a_{2,2}$	$a_{0,8} = \frac{7\eta}{10}a_{1,5}^2$
8	$a_{0,7} = 7a_{1,3}a_{1,5} + \frac{35}{3}a_{1,3}^2a_{2,3} - \frac{35}{9}a_{1,3}^3a_{3,1}$ $- \frac{35}{8}(a_{1,4} - 2a_{1,3}a_{2,2})^2$	$a_{0,10} = 3\eta a_{1,5}a_{1,7} + \frac{21}{16}a_{1,5}^2a_{2,4}$

(6.14)

Comment to Table (6.14). In this table, the variables are denoted by $x = k_1$, $y = k_2$, and $z = (k_3, \dots, k_\nu)$, moreover $f_{x^a y^b z^c} := \frac{\partial^{a+b+c} f(0)}{\partial x^a \partial y^b \partial z^c}$. Furthermore, one denotes $a_{0,j} = a_{j-1}$ where a_μ are the real numbers assigned according to Statement 6.1 to the function $f(0, k_2, \dots, k_\nu)$, see Table (6.5). Other real numbers $a_{i,j} = \frac{\partial f^{i+j}(0)}{\partial x^i \partial y^j} + \dots$ in Table (6.14) are similar to the numbers $a_{0,j}$ and are defined by the formulae

$$\begin{aligned}
a_{1,3} &= f_{xyz} - 3f_{xyz}f_{z^2}^{-1}f_{y^2z}, \quad a_{2,2} = f_{x^2y^2} - 2f_{z^2}^{-1}[f_{xyz}]^2 - f_{x^2z}f_{z^2}^{-1}f_{y^2z}, \\
a_{1,4} &= f_{xy^4} - 6f_{xy^2z}f_{z^2}^{-1}f_{y^2z} - 4f_{y^3z}f_{z^2}^{-1}f_{xyz} + 3f_{xz^2}[f_{z^2}^{-1}f_{y^2z}]^2 + 12f_{y^2z}f_{z^2}^{-1}f_{y^2z}f_{z^2}^{-1}f_{xyz}, \\
a_{2,3} &= f_{x^2y^3} - f_{x^2z}f_{z^2}^{-1}f_{y^3z} - 3(f_{x^2yz} - f_{x^2z}f_{z^2}^{-1}f_{yz^2} - 2f_{xyz}f_{z^2}^{-1}f_{xz^2})f_{z^2}^{-1}f_{y^2z} \\
&\quad - 6(f_{xyz}f_{z^2}^{-1}f_{xy^2z} - f_{yz^2}[f_{z^2}^{-1}f_{xyz}]^2), \\
a_{1,5} &= f_{xy^5} - 10f_{y^3z}f_{z^2}^{-1}f_{xy^2z} - 30f_{xz^2}f_{z^2}^{-1}f_{yz^2}[f_{z^2}^{-1}f_{y^2z}]^2 - 60f_{xyz}f_{z^2}^{-1}[f_{yz^2}f_{z^2}^{-1}]^2f_{y^2z} - 5f_{y^4z}f_{z^2}^{-1}f_{xyz} \\
&\quad - 10f_{xy^3z}f_{z^2}^{-1}f_{y^2z} - 15f_{z^3}[f_{z^2}^{-1}f_{xyz}]^3 + 15f_{xyz}f_{z^2}^{-1}[f_{z^2}^{-1}f_{y^2z}]^2 + 30f_{xyz}f_{z^2}^{-1}f_{y^2z}f_{z^2}^{-1}f_{y^2z} \\
&\quad + 10f_{y^2z}f_{z^2}^{-1}f_{xz^2}f_{z^2}^{-1}f_{y^3z} + 20f_{xyz}f_{z^2}^{-1}f_{yz^2}f_{z^2}^{-1}f_{y^3z} + 30f_{y^2z}f_{z^2}^{-1}f_{yz^2}f_{z^2}^{-1}f_{xy^2z}.
\end{aligned}$$

Moreover, if f is an even function then

$$\begin{aligned}
a_{2,4} &= f_{x^2y^4} - 4f_{x^2yz}f_{z^2}^{-1}f_{y^3z} - 6f_{z^2}^{-1}[f_{xy^2z}]^2, \\
a_{1,7} &= f_{xy^7} - 35f_{xy^4z}f_{z^2}^{-1}f_{y^3z} - 21f_{xy^2z}f_{z^2}^{-1}(f_{y^5z} - 10[f_{y^2z}f_{z^2}^{-1}]^2f_{y^3z}) + 70f_{xyz}f_{z^2}^{-1}[f_{y^3z}]^2.
\end{aligned}$$

One can obtain $a_{j,i}$ from $a_{i,j}$ by replacing all partial derivatives with respect to x by partial derivatives with respect to y , and vica-versa. The above formulae are written for $\nu = 3$, but they are easily transformed to the corresponding formulae for any $\nu \geq 2$, see (6.6). In the right column of Table (6.14), the function f is supposed to be even. This leads to a simplification of the formulae for the numbers $a_{i,j}$.

One proves Statements 6.7 and 6.8 similarly to the proof of Statement 6.1, using the Newton ruler method [1, v. 1, 12.6], see also 3.12. Namely, due to the parametric Morse lemma for arbitrary smooth functions (analogous to Theorem 3.6), in a small neighbourhood of the origin, there exists a smooth change of variables $(k_1, k_2, k_3, \dots, k_\nu) \rightarrow (k_1, k_2, \tilde{k}_3, \dots, \tilde{k}_\nu)$ leaving the origin fixed and reducing f to the form $f = a + \psi(k_1, k_2) + Q(\tilde{k}_3, \dots, \tilde{k}_\nu)$, where a is a constant, Q is a nondegenerate quadratic form, and $\psi(k_1, k_2) = o(k_1^2 + k_2^2)$ as $(k_1, k_2) \rightarrow 0$. Let $P(k_1, k_2) = Ak_1^3 + Bk_1^2k_2 + Ck_1k_2^2 + Dk_2^3$ be the Taylor polynomial of degree 3 of the function ψ at zero. It follows from the Newton ruler method that the germ at the origin of the function ψ in two variables has a singularity of type D_4 if and only if three lines, which are defined by the linear factors of the decomposition of P , are pairwise different, compare 3.12. This is equivalent to the fact that the discriminant Δ_3 of the polynomial $Ax^3 + Bx^2 + Cx + D$ does not vanish [11]. To see the validity of the formulae for the coefficients A, B, C, D in Statement 6.7, one uses the fact that the definition of the «cubical»

part (terms of degree ≤ 3) of a function at a critical point is well defined [1], and the fact that the coordinate plane Ok_1k_2 coincides with the kernel of the second differential of f at zero, both for the initial variables and for the new variables. Statement 6.7 is thus proved.

In order to prove Statement 6.8, we observe that $A = C = 0$ and $B = \frac{1}{2}$. If $D \neq 0$ then $\Delta_3 \neq 0$ and, by the above arguments, f has a singularity at the origin of type D_4 . Now we assume that $D = 0$ and $\mu \geq 5$. Following the method of ruler turning, we perform the change of variables $(k_1, k_2) \rightarrow (\tilde{k}_1, \tilde{k}_2)$ given by $\tilde{k}_1 = k_1 + \lambda_2 k_2^2 + \dots + \lambda_{[\mu/2]-1} k_2^{[\mu/2]-1}$, $\tilde{k}_2 = k_2$, where the λ_j are determined implicitly (also uniquely and independently of μ) by the following condition: after this change of variables the coefficients at the terms $\tilde{k}_1 \tilde{k}_2^j$, $3 \leq j \leq [\mu/2]$, of the Taylor series of ψ in $\tilde{k}_1, \tilde{k}_2$, centred at the origin, vanish. Due to the ruler method, the function ψ (and, hence, the function f) has a singularity at the origin of type D_μ if and only if the coefficient at the term $\tilde{k}_2^j$ in the Taylor series of ψ at zero vanishes for any $j \leq \mu - 2$ and does not vanish for $j = \mu - 1$.

Therefore, it remains to express the numbers $d_\mu := \frac{\partial^{\mu-1} \psi(0)}{\partial k_2^{\mu-1}}$ in terms of the values of the partial derivatives of f at the origin. First we will express the numbers $a_{q,p} := \frac{\partial^{q+p} \psi(0)}{\partial k_1^q \partial k_2^p} = \frac{\partial^{q+p} f(0)}{\partial k_1^q \partial k_2^p} + \dots$. Computations, analogous to those for the numbers $a_\mu = \psi^{(\mu+1)}(0) = \frac{\partial^{\mu+1} f(0)}{\partial k_1^{\mu+1}} + \dots$ (see the proof of Statement 6.1), lead to the above formulas for $a_{q,p}$, see the comment to Table (6.14). By means of the method of ruler turning, one easily obtains an expression for the coefficient λ_j of the change of variables $(k_1, k_2) \rightarrow (\tilde{k}_1, \tilde{k}_2)$ in terms of the numbers $a_{q,p}$, $q \geq 1$, $2q + p \leq j + 3$. After performing this change of variables, one obtains the required expression $d_\mu = a_{0,\mu-1} + \dots$ in terms of the numbers $a_{q,p}$, $2q + p \leq \mu - 1$. This proves Statement 6.8.

Statements 6.10 and 6.12 are similarly proved based on Theorem 3.6 and the results of Section 3.12, see the cases 1 and 2.

Statements 6.1, 6.3, 6.7, 6.8, 6.10, 6.12, and Theorem 3.9 demonstrate that singularities of type A_μ , D_μ , and even singularities of type $A_{e,\mu}$ have codimension $c = \mu - 1$, while even singularities of class $X_{e,\mu}^{\pm\pm}$ have even codimension $c = \mu - 2$. That is, these singularities are non-removable for typical families with $l \geq c$ parameters on the entire family. Besides, it follows from these assertions that the only singularities which appear in typical families with $l = 1, 2, 3, 4$ parameters are A_1 , A_2 , A_3 , A_4 , A_5 , D_4 , and D_5 (the seven of Thom). In additional, in the case of typical even families, the only even singularities which appear in such families are $A_{e,1}$, $A_{e,2}$, $A_{e,3}$, $A_{e,4}$, $A_{e,5}$, $X_{e,5}$, and $X_{e,6}$.

Here, by *typical* (even) families of functions, we understand (even) families which form an open dense subspace in the space of all (even) families of functions (in the convergence topology with a finite number of derivatives on each compact set).

Actually, due to Tables (2.3) and (2.7), in the case of five parameters, three more singularities A_6 , D_6 , and E_6 , as well as five more even singularities $A_{e,6}$, $X_{e,7}$, $Y_{e,3,3}$, $\tilde{Y}_{e,3}$, and $Z_{e,7}$ appear.

7 Conditions for versal deformations

In this section, we describe conditions on a deformation of a germ to be versal or, equivalently, stable, see 4.9.

It is known that the versal property is fulfilled for *typical* deformations with $l \geq \mu - 1$ parameters, i.e. deformations which form an open dense subspace in the space of all l -parameter deformations of a given germ (in the convergence topology with a finite number of derivatives on each compact set). However, the deformations which appear in practice are often not arbitrary, but have a special form. Therefore, in addition to the question about the determination of the type of a singular point of a given function $f(k)$, the following questions are also of interest for applications: whether an l -parameter deformation $F(k, \lambda)$ of the germ of this function is versal, and which normal form this deformation has near the singular point.

In the case of singularities of codimension 0 (Morse functions and even Morse functions), answers are given by the parametric Morse lemma for arbitrary functions and even functions. The following assertions give conditions on l -parameter deformations with $l = 1, 2, 3, 4$ parameters to be versal for «usual» singularities and even singularities. As we have mentioned above, these assertions can be considered as analogues of the parametric Morse lemma for degenerate singularities.

In the following assertions, one considers a family of functions $F = F(k, \lambda)$ in a variable $k = (k_1, \dots, k_\nu)$ with a parameter $\lambda = (\lambda_1, \dots, \lambda_l)$. One denotes by $\frac{\partial^{i_1+\dots+i_\nu+1} F(0,0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu} \partial \lambda}$ the l -dimensional vector (i.e. the element of $\mathbb{R}^l$) with components $\frac{\partial^{i_1+\dots+i_\nu+1} F(0,0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu} \partial \lambda_j}$, $1 \leq j \leq l$.

Statement 7.1 (A deformation of a singularity A_μ) *Let, under the hypothesis of Statement 6.1, $F : \mathbb{R}^\nu \times \mathbb{R}^l \rightarrow \mathbb{R}$ be a smooth l -parameter deformation of the germ of f at the origin, i.e. $f = F(\cdot, 0)$. Then there exists a sequence of vectors $\bar{a}_\mu \in \mathbb{R}^l$, $\mu = 2, 3, \dots$, depending on partial derivatives of f at the origin, and possessing the following properties:*

1. $\bar{a}_\mu = \frac{\partial^\mu F(0,0)}{\partial k_1^{\mu-1} \partial \lambda} + \dots$, where the additional terms form a linear combination of vectors $\frac{\partial^{i_1+\dots+i_\nu+1} F(0,0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu} \partial \lambda}$, $i_1 + \dots + i_\nu + 1 < \mu$, $(i_2, \dots, i_\nu) \neq (0, \dots, 0)$, the coefficients of which are polynomials in the values of the partial derivatives of order $\leq \mu$ of f at the origin (except the derivatives with respect to the variable k_1 only) and the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 2, \dots, \nu$. These polynomials have integer coefficients and vanishing free terms. The vectors $\bar{a}_\mu$ are given in Table (7.3) for $\mu = 2, 3, 4, 5$.

2. Suppose that the conditions (6.2) are fulfilled, i.e. the germ of f at the origin has a singularity of type A_μ . The deformation F is R^+ -versal if and only if the vectors

$$\bar{a}_2, \dots, \bar{a}_\mu$$

form a linearly independent system in $\mathbb{R}^l$ (in particular, $l \geq \mu - 1$).

Moreover, if (6.2) holds and the above vectors form a linearly independent system in $\mathbb{R}^l$ then F reduces to the form $F = a(\tilde{\lambda}) + a_\mu \tilde{k}_1^{\mu+1} + Q(\tilde{k}_2, \dots, \tilde{k}_\nu) + \tilde{\lambda}_1 \tilde{k}_1 + \dots + \tilde{\lambda}_{\mu-1} \tilde{k}_1^{\mu-1}$ in some neighbourhood of the origin by means of a regular change of variables $(k, \lambda) \rightarrow (\tilde{k}, \tilde{\lambda})$ leaving the origin fixed and having the form $\tilde{k} = \tilde{k}(k, \lambda)$, $\tilde{\lambda} = \tilde{\lambda}(\lambda)$. Here a is smooth function, and Q is a nondegenerate quadratic form in $\nu - 1$ variables.

Statement 7.2 (An even deformation of a singularity $A_{e,\mu}$) *Let, under the hypothesis of Statement 6.3, $f(0) = 0$ and let $F : \mathbb{R}^\nu \times \mathbb{R}^l \rightarrow \mathbb{R}$ be a smooth l -parameter even deformation of the germ of f at the origin. That is, $f = F(\cdot, 0)$ and, for each λ , the function $F(\cdot, \lambda)$ is even and has a vanishing value at the origin. Then there exists a sequence of vectors $\bar{a}_{e,\mu} \in \mathbb{R}^l$, $\mu = 2, 3, \dots$, depending on partial derivatives of F at the origin, and possessing the following properties:*

1. $\bar{a}_{e,\mu} = \bar{a}_{2\mu-1} = \frac{\partial^{2\mu-1} F(0,0)}{\partial k_1^{2\mu-2} \partial \lambda} + \dots$, where the vectors $\bar{a}_{2\mu-1}$ are assigned according to Statement 7.1 to the even deformation F , which is considered as a usual deformation. The vectors $\bar{a}_{e,\mu}$ are given in Table (7.3) for $\mu = 2, 3, 4$.

2. Suppose that the conditions (6.4) are fulfilled, i.e. the even germ of f at the origin has a singularity of type $A_{e,\mu}$. Then the even deformation F is R_O -versal if and only if the vectors

$$\bar{a}_{e,2}, \dots, \bar{a}_{e,\mu}$$

form a linearly independent system in $\mathbb{R}^l$ (in particular, $l \geq \mu - 1$).

Moreover, if (6.4) holds and the above vectors form a linearly independent system in $\mathbb{R}^l$ then F reduces to the form $F = a_{e,\mu} \tilde{k}_1^{2\mu} + Q(\tilde{k}_2, \dots, \tilde{k}_\nu) + \tilde{\lambda}_1 \tilde{k}_1^2 + \dots + \tilde{\lambda}_{\mu-1} \tilde{k}_1^{2(\mu-1)}$ in some neighbourhood of the origin by means of a regular change of variables $(k, \lambda) \rightarrow (\tilde{k}, \tilde{\lambda})$ leaving the origin fixed and having the form $\tilde{k} = \tilde{k}(k, \lambda)$, $\tilde{\lambda} = \tilde{\lambda}(\lambda)$ with $\tilde{k}(-k, \lambda) = -\tilde{k}(k, \lambda)$. Here Q is as in Statement 7.1.

μ	$\bar{a}_\mu$	$\bar{a}_{e,\mu}$
2	$F_{y\lambda}$	$F_{y^2\lambda}$
3	$F_{y^2\lambda} - f_{y^2z}f_{z^2}^{-1}F_{z\lambda}$	$F_{y^4\lambda} - 4f_{y^3z}f_{z^2}^{-1}F_{yz\lambda}$
4	$F_{y^3\lambda} - (f_{y^3z} - 3f_{y^2z}f_{z^2}^{-1}f_{yz^2})f_{y^2}^{-1}F_{z\lambda}$ $- 3f_{y^2z}f_{z^2}^{-1}F_{yz\lambda}$	$F_{y^6\lambda} - 6(f_{y^5z} - 10f_{y^3z}f_{z^2}^{-1}f_{yz^2})f_{z^2}^{-1}F_{yz\lambda}$ $- 20f_{y^3z}f_{z^2}^{-1}F_{yz\lambda} + 4F_{z^2\lambda}[f_{z^2}^{-1}f_{y^3z}]^2$
5	$F_{y^4\lambda} - 4(f_{y^3z} - 3f_{y^2z}f_{z^2}^{-1}f_{yz^2})f_{z^2}^{-1}F_{yz\lambda}$ $- 6f_{y^2z}f_{z^2}^{-1}F_{yz\lambda} + F_{z^2\lambda}[f_{z^2}^{-1}f_{y^2z}]^2$ $- (f_{y^4z} - 4f_{y^3z}f_{z^2}^{-1}f_{yz^2} + 9f_{y^2z}[f_{z^2}^{-1}f_{yz^2}]^2$ $- 6f_{y^2z}f_{z^2}^{-1}f_{yz^2} - f_{z^3}[f_{z^2}^{-1}f_{y^2z}]^2)f_{z^2}^{-1}F_{z\lambda}$	$F_{y^8\lambda} + \dots$

(7.3)

Comment to Table (7.3). In this table, the variables and the parameters are denoted by $y = k_1$, $z = (k_2, \dots, k_\nu)$, and $\lambda = (\lambda_1, \dots, \lambda_l)$, resp., moreover $f_{y^a z^b} := \frac{\partial^{a+b} f(0)}{\partial y^a \partial z^b}$ and $F_{y^a z^b \lambda} := \frac{\partial^{a+b+1} F(0,0)}{\partial y^a \partial z^b \partial \lambda}$. The formulae in the table are written for $\nu = 2$, $l = 1$, but they are easily transformed to the corresponding formulae for any $\nu, l \geq 1$, see (6.6).

For a proof of Statement 7.1, let us perform, for any sufficiently small value of the parameter λ , a change of variables similar to that in the proof of Statement 6.1. In the new variables $k_1, \tilde{K}_2, \dots, \tilde{K}_\nu, \lambda$, we have $F = A(\lambda) + \Psi(k_1, \lambda) + Q(\tilde{K}_2, \dots, \tilde{K}_\nu)$, where A and Ψ are smooth functions, $A(0) = a$, $\Psi(\cdot, 0) = \psi$, $\tilde{K} = \tilde{K}(k, \lambda)$. Since the function $\psi(k_1)$ has order $\mu + 1$ at the origin, the germ of the function k_1^μ belongs to the Jacobian ideal $I_{\nabla\psi}$. It follows that the germs of the functions $k_1, \dots, k_1^{\mu-1}$ are generators of the local algebra $Q_{\nabla\psi}$ of the gradient map of the function ψ at the origin (considered as a vector space), and that the differential operators $\frac{\partial}{\partial k_1}|_0, \frac{\partial^2}{\partial k_1^2}|_0, \dots, \frac{\partial^{\mu-1}}{\partial k_1^{\mu-1}}|_0$ are generators of its dual space $Q_{\nabla\psi}^*$. It follows from this, from the conditions on a deformation to be infinitesimally R^+ -versal, and from the versality theorem (see the theorems from [1, v. 1, 8.2 and 8.3]), that a deformation Ψ of a function ψ is R^+ -versal if and only if the vectors $\bar{a}_i := \frac{\partial^i \Psi(0,0)}{\partial k_1^{i-1} \partial \lambda} \in \mathbb{R}^l$, $i = 2, \dots, \mu$ are linearly independent.

In order to compute the vectors $\bar{a}_\mu$, observe that $\Psi(k_1, \lambda) = F(k_1, K_2(k_1, \lambda), \dots, K_\nu(k_1, \lambda); \lambda)$ where $K_i = K_i(k_1, \lambda)$, $i = 2, \dots, \nu$ are the functions which define the coordinate plane $Ok_1\lambda$ of the coordinate system $k_1, \tilde{K}_2, \dots, \tilde{K}_\nu, \lambda$ and, therefore, are implicitly defined by the system of equations

$$\frac{\partial F}{\partial k_i}(k_1, K_2(k_1, \lambda), \dots, K_\nu(k_1, \lambda); \lambda) = 0, \quad i = 2, \dots, \nu.$$

In particular, $K_i(k_1, 0) = k_i(k_1)$, $i = 2, \dots, \nu$, see the proof of Statement 6.1. One now sets to zero the value of each composite partial derivative of bi-order $(m, 1)$, $m \leq \mu - 2$, at the origin with respect to (k_1, λ) on the left-hand sides of these equations. This will give expressions for composite partial derivatives at the origin of bi-order $(m, 1)$ with respect to (k_1, λ) of the functions $K_i(k_1, \lambda)$. By substituting these expressions and the expressions for $k_i'(0), \dots, k_i^{(\mu)}(0)$ into the composite partial derivative of order $(\mu - 1, 1)$ of the composite function Ψ at the origin with respect to (k_1, λ) , we obtain the required expression for $\bar{a}_\mu$.

Statement 7.2 is similarly proved on the basis of Theorem 4.6.

Statement 7.4 (A deformation of a singularity D_μ) *Let, under the hypothesis of Statement 6.8, $F : \mathbb{R}^\nu \times \mathbb{R}^l \rightarrow \mathbb{R}$ be a smooth l -parameter deformation of the germ of f at the origin, i.e. $f = F(\cdot, 0)$. Then there exists a sequence of vectors $\bar{d}_\mu \in \mathbb{R}^l$, $\mu = 1, 2, 3, \dots$, depending on partial derivatives of F at the origin, and possessing the following properties:*

1. $\bar{d}_2 = \frac{\partial^2 F(0,0)}{\partial k_1 \partial \lambda}$ and $\bar{d}_1 = \frac{\partial^3 F(0,0)}{\partial k_1^2 \partial \lambda} + \dots$ are equal to the vectors $\bar{a}_2$ and $\bar{a}_3$, resp., which are assigned according to Statement 7.1 to the deformation $F(k_1, 0, k_3, \dots, k_\nu; \lambda)$ of the germ of

$f(k_1, 0, k_3, \dots, k_\nu)$ at the origin. The vectors $\bar{d}_3 = \frac{\partial^2 F(0,0)}{\partial k_2 \partial \lambda}$ and $\bar{d}_4 = \frac{\partial^3 F(0,0)}{\partial k_2^2 \partial \lambda} + \dots$ are similarly defined by means of the deformation $F(0, k_2, k_3, \dots, k_\nu; \lambda)$ of the germ of $f(0, k_2, k_3, \dots, k_\nu)$ at the origin. For $\mu \geq 5$, one has $\bar{d}_\mu = \frac{\partial^{\mu-1} F(0,0)}{\partial k_2^{\mu-2} \partial \lambda} + \dots$, where the additional terms form a linear combination of vectors $\frac{\partial^{i_1+\dots+i_\nu+1} F(0,0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu} \partial \lambda}$, $i_1 + \dots + i_\nu + 1 < \mu - 1$, $(i_1, i_3, \dots, i_\nu) \neq (0, \dots, 0)$, the coefficients of which are polynomials in the values of the partial derivatives of order $\leq \mu - 1$ of f at the origin (except the derivatives with respect to the variable k_2 only) and the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 3, \dots, \nu$. These polynomials have rational coefficients and vanishing free terms. The vectors $\bar{d}_\mu$ are given in Table (7.6) for $\mu = 1, 2, 3, 4, 5, 6, 7$.

2. Suppose that the conditions (6.9) are fulfilled, i.e. the germ of f at the origin has a singularity of type $D_\mu^\pm$, $\mu \geq 4$. The deformation F is R^+ -versal if and only if the vectors

$$\bar{d}_2, \dots, \bar{d}_{\mu-1}, \bar{d}_\mu - d_\mu \bar{d}_1$$

form a linearly independent system in $\mathbb{R}^l$ (in particular, $l \geq \mu - 1$).

Moreover, if (6.9) holds and the above vectors form a linearly independent system in $\mathbb{R}^l$ then F reduces to the form $F = d(\tilde{\lambda}) + \tilde{k}_1^2 \tilde{k}_2 + d_\mu \tilde{k}_2^{\mu-1} + Q(\tilde{k}_3, \dots, \tilde{k}_\nu) + \tilde{\lambda}_1 \tilde{k}_1 + \tilde{\lambda}_2 \tilde{k}_1^2 + \tilde{\lambda}_3 \tilde{k}_2 + \tilde{\lambda}_4 \tilde{k}_2^2 + \dots + \tilde{\lambda}_{\mu-1} \tilde{k}_2^{\mu-3}$ in some neighbourhood of the origin by means of a regular change of variables $(k, \lambda) \rightarrow (\tilde{k}, \tilde{\lambda})$ leaving the origin fixed and having the form $\tilde{k} = \tilde{k}(k, \lambda)$, $\tilde{\lambda} = \tilde{\lambda}(\lambda)$. Here d is a smooth function, and Q is a nondegenerate quadratic form in $\nu - 2$ variables.

Statement 7.5 (An even deformation of a singularity $X_{e,\mu}$) Let, under the hypothesis of Statement 6.12, $f(0) = 0$ and let $F : \mathbb{R}^\nu \times \mathbb{R}^l \rightarrow \mathbb{R}$ be a smooth l -parameter even deformation of the germ of f at the origin. That is, $f = F(\cdot, 0)$ and, for each λ , the function $F(\cdot, \lambda)$ is even and has a vanishing value at the origin. Then there exists a sequence of vectors $\bar{x}_{e,\mu} = \bar{x}_{e,\mu}^{\varepsilon,\eta} \in \mathbb{R}^l$, $\mu = 1, 2, 3, \dots$ depending on partial derivatives of F at the origin and possessing the following properties:

1. $\bar{x}_{e,2}^{\varepsilon,\eta} = \frac{\partial^3 F(0,0)}{\partial k_1^2 \partial \lambda}$, $\bar{x}_{e,3}^{\varepsilon,\eta} = \frac{\partial^3 F(0,0)}{\partial k_1 \partial k_2 \partial \lambda}$, $\bar{x}_{e,1}^{\varepsilon,\eta} = \varepsilon(\frac{\partial^5 F(0,0)}{\partial k_1^4 \partial \lambda} + \dots) - 12\eta(\frac{\partial^5 F(0,0)}{\partial k_1^2 \partial k_2^2 \partial \lambda} + \dots)$, see Table (7.6).

For $\mu \geq 4$, one has $\bar{x}_{e,\mu}^{\varepsilon,\eta} = \frac{\partial^{2\mu-5} F(0,0)}{\partial k_2^{2\mu-6} \partial \lambda} + \dots$, where the additional terms form a linear combination of vectors $\frac{\partial^{i_1+\dots+i_\nu+1} F(0,0)}{\partial k_1^{i_1} \dots \partial k_\nu^{i_\nu} \partial \lambda}$, $i_1 + \dots + i_\nu + 1 < 2\mu - 5$, $(i_1, i_3, \dots, i_\nu) \neq (0, \dots, 0)$, the coefficients of which are polynomials in the values of the partial derivatives of order $\leq 2\mu - 4$ of f at the origin (except the derivatives with respect to the variable k_2 only) and the components of the inverse of the matrix $\|\frac{\partial^2 f(0)}{\partial k_i \partial k_j}\|$, $i, j = 3, \dots, \nu$. These polynomials have rational coefficients and vanishing free terms. The vectors $\bar{x}_{e,\mu}^{\varepsilon,\eta}$ are given in Table (7.6) for $\mu = 1, 2, 3, 4, 5, 6$.

2. Suppose that the conditions (6.13) are fulfilled, i.e. the even germ of f at the origin has a singularity of class $X_{e,\mu}^{\varepsilon,\eta}$, $\mu \geq 5$. The even deformation F is R_O -versal if and only if the vectors

$$\bar{x}_{e,2}^{\varepsilon,\eta}, \dots, \bar{x}_{e,\mu-1}^{\varepsilon,\eta}, \bar{x}_{e,\mu}^{\varepsilon,\eta} + \frac{\mu-3}{48} x_{e,\mu}^{\varepsilon,\eta} \bar{x}_{e,1}^{\varepsilon,\eta}$$

form a linearly independent system in $\mathbb{R}^l$ (in particular, $l \geq \mu - 1$).

Moreover, if (6.13) holds and the above vectors form a linearly independent system in $\mathbb{R}^l$, then F reduces to the form $F = \varepsilon \tilde{k}_1^4 + \eta \tilde{k}_1^2 \tilde{k}_2^2 + \frac{x_{e,\mu}^{\varepsilon,\eta}}{(2\mu-6)!} \tilde{k}_2^{2\mu-6} + Q(\tilde{k}_3, \dots, \tilde{k}_\nu) + \tilde{\lambda}_1 \tilde{k}_1^2 + \tilde{\lambda}_2 \tilde{k}_1 \tilde{k}_2 + \tilde{\lambda}_3 \tilde{k}_2^2 + \tilde{\lambda}_4 \tilde{k}_1^4 + \tilde{\lambda}_5 \tilde{k}_2^4 + \dots + \tilde{\lambda}_{\mu-1} \tilde{k}_2^{2\mu-8}$ in some neighbourhood of the origin by means of a regular change of variables $(k, \lambda) \rightarrow (\tilde{k}, \tilde{\lambda})$ leaving the origin fixed and having the form $\tilde{k} = \tilde{k}(k, \lambda)$, $\tilde{\lambda} = \tilde{\lambda}(\lambda)$ with $\tilde{k}(-k, \lambda) = -\tilde{k}(k, \lambda)$. Here Q is as in Statement 7.4.

μ	$\bar{d}_\mu$	$\bar{x}_{e,\mu} = \bar{x}_{e,\mu}^{\varepsilon,\eta}$
1	$\bar{a}_{2,0} = F_{x^2\lambda} - f_{x^2z}f_{z^2}^{-1}F_{z\lambda}$	$\varepsilon\bar{a}_{4,0} - 12\eta\bar{a}_{2,2}$
2	$\bar{a}_{1,0} = F_{x\lambda}$	$\bar{a}_{2,0} = F_{x^2\lambda}$
3	$\bar{a}_{0,1} = F_{y\lambda}$	$\bar{a}_{1,1} = F_{xy\lambda}$
4	$\bar{a}_{0,2} = F_{y^2\lambda} - f_{y^2z}f_{z^2}^{-1}F_{z\lambda}$	$\bar{a}_{0,2} = F_{y^2\lambda}$
5	$\bar{a}_{0,3} - 3a_{1,3}\bar{a}_{1,1}$	$\bar{a}_{0,4} - \frac{\eta}{10}a_{1,5}\bar{a}_{1,1}$
6	$\bar{a}_{0,4} - 2a_{1,3}\bar{a}_{1,2} - (a_{1,4} - 2a_{2,2}a_{1,3})\bar{a}_{1,1} + \frac{1}{3}a_{1,3}^2\bar{a}_{2,0}$	$\bar{a}_{0,6} - \frac{\eta}{2}a_{1,5}\bar{a}_{1,3} + \frac{1}{160}a_{1,5}^2\bar{a}_{2,0}$ $+ (\frac{1}{16}a_{1,5}a_{2,4} - \frac{1}{14}\eta a_{1,7})\bar{a}_{1,1}$
7	$\bar{a}_{0,5} - \frac{10}{3}a_{1,3}\bar{a}_{1,3} + \frac{5}{3}a_{1,3}^2\bar{a}_{2,1}$ $-(a_{1,5} - \frac{10}{3}a_{2,3}a_{1,3} + \frac{5}{3}a_{3,1}a_{1,3}^2)\bar{a}_{1,1}$ $+\frac{5}{6}(a_{1,4} - 2a_{2,2}a_{1,3})(a_{1,3}\bar{a}_{2,0} - 3\bar{a}_{1,2} - 3a_{2,2}\bar{a}_{1,1})$	$F_{y^8\lambda} + \dots$

(7.6)

Comment to Table (7.6). In this table, the variables and the parameters are denoted by $x = k_1$, $y = k_2$, $z = (k_3, \dots, k_\nu)$, and $\lambda = (\lambda_1, \dots, \lambda_l)$, resp., moreover $f_{x^a y^b z^c} := \frac{\partial^{a+b+c} f(0)}{\partial x^a \partial y^b \partial z^c}$ and $F_{x^a y^b z^c \lambda} := \frac{\partial^{a+b+c+1} F(0,0)}{\partial x^a \partial y^b \partial z^c \partial \lambda}$. The real numbers $a_{i,j} = \frac{\partial f^{i+j}(0)}{\partial x^i \partial y^j} + \dots$ are the same as those in Table (6.14). Furthermore, one denotes $\bar{a}_{0,j} = \bar{a}_{j+1}$ where $\bar{a}_\mu \in \mathbb{R}^l$ are the vectors assigned according to Statement 7.1 to the deformation $F(0, k_2, \dots, k_\nu, \lambda)$ of $f(0, k_2, \dots, k_\nu)$, see Table (7.3). Other vectors $\bar{a}_{i,j} = \frac{\partial F^{i+j+1}(0,0)}{\partial x^i \partial y^j \partial \lambda} + \dots \in \mathbb{R}^l$ in Table (7.6) are similar to the vectors $\bar{a}_{0,j}$ and are defined by the formulae

$$\begin{aligned}\bar{a}_{1,1} &= F_{xy\lambda} - f_{xyz}f_{z^2}^{-1}F_{z\lambda}, \\ \bar{a}_{1,2} &= F_{xy^2\lambda} - f_{y^2z}f_{z^2}^{-1}F_{xz\lambda} - 2f_{xyz}f_{z^2}^{-1}F_{yz\lambda} - (f_{xy^2z} - f_{y^2z}f_{z^2}^{-1}f_{xz^2} - 2f_{xyz}f_{z^2}^{-1}f_{yz^2})f_{z^2}^{-1}F_{z\lambda}, \\ \bar{a}_{1,3} &= F_{xy^3\lambda} - (f_{y^3z} - 3f_{y^2z}f_{z^2}^{-1}f_{yz^2})f_{z^2}^{-1}F_{xz\lambda} - 3(f_{xy^2z} - f_{y^2z}f_{z^2}^{-1}f_{xz^2} - 2f_{xyz}f_{z^2}^{-1}f_{yz^2})f_{z^2}^{-1}F_{yz\lambda} \\ &\quad - 3f_{xyz}f_{z^2}^{-1}F_{y^2z\lambda} - 3f_{y^2z}f_{z^2}^{-1}(F_{xyz\lambda} - F_{z^2\lambda}f_{z^2}^{-1}f_{xyz}) - (f_{xy^3z} - f_{y^3z}f_{z^2}^{-1}f_{xz^2} - 3f_{xy^2z}f_{z^2}^{-1}f_{yz^2} \\ &\quad - 3(f_{xyz}(f_{z^2}^{-1}f_{yz^2} - 2[f_{z^2}^{-1}f_{yz^2}]^2) + f_{y^2z}f_{z^2}^{-1}(f_{xyz^2} - 2f_{yz^2}f_{z^2}^{-1}f_{xz^2} - f_{z^3}f_{z^2}^{-1}f_{xyz})))f_{z^2}^{-1}F_{z\lambda}.\end{aligned}$$

Moreover, if $F(\cdot, \lambda)$ is an even function for any λ then

$$\bar{a}_{2,2} = F_{x^2y^2\lambda} - 2f_{xy^2z}f_{z^2}^{-1}F_{xz\lambda} - 2f_{x^2yz}f_{z^2}^{-1}F_{yz\lambda}.$$

One can obtain $\bar{a}_{j,i}$ from $\bar{a}_{i,j}$ by replacing all partial derivatives with respect to x by partial derivatives with respect to y , and vica-versa. The above formulae are written for $\nu = 3$, $l = 1$, but they are easily transformed to the corresponding formulae for any $\nu \geq 2$, $l \geq 1$, see (6.6). In the right column of Table (7.6), the function $F(\cdot, \lambda)$ is supposed to be even for each λ . This leads to a simplification of the formulae for the vectors $\bar{a}_{i,j}$.

For a proof of Statement 7.4, let us perform, for any sufficiently small value of the parameter λ , a change of variables similar to that in the proof of Statement 6.8. In the new variables $k_1, k_2, \tilde{K}_3, \dots, \tilde{K}_\nu, \lambda$, we have $F = A(\lambda) + \Psi(k_1, k_2, \lambda) + Q(\tilde{K}_3, \dots, \tilde{K}_\nu)$, where A and Ψ are smooth functions, $A(0) = a$, $\Psi(\cdot, 0) = \psi$, $\tilde{K} = \tilde{K}(k, \lambda)$. Now let us perform the change of variables $(k_1, k_2) \rightarrow (\tilde{k}_1, \tilde{k}_2)$ given by $\tilde{k}_1 = k_1 + \lambda_2 k_2^2 + \dots + \lambda_{\mu-3} k_2^{\mu-3}$, $\tilde{k}_2 = k_2$, where the λ_j are determined implicitly (also uniquely and independently of μ) by the following condition: after this change of variables the coefficients at the terms $\tilde{k}_1 \tilde{k}_2^j$, $3 \leq j \leq \mu - 2$, of the Taylor series of ψ in $\tilde{k}_1, \tilde{k}_2$, centred at the origin, vanish, see the proof of Statement 6.8. Denote by $\tilde{\psi}(\tilde{k}_1, \tilde{k}_2)$ and $\tilde{\Psi}(\tilde{k}_1, \tilde{k}_2, \lambda)$ the functions which are obtained from the functions $\psi(k_1, k_2)$ and $\Psi(k_1, k_2, \lambda)$, resp., after this change of variables. Thus, the coefficients at the terms $\tilde{k}_1 \tilde{k}_2^j$, $j \leq \mu - 2$, of the Taylor series of $\tilde{\psi}$, centred at zero, vanish. It follows from the proof of Statement 6.8 that, for each $j \leq 2\mu - 4$, the coefficient at

the term $\tilde{k}_2^j$ equals $\frac{d_{j+1}}{j!}$, therefore the coefficients at the terms $\tilde{k}_2^j$, $j \leq \mu - 2$, also vanish, and the coefficient at the term $\tilde{k}_2^{\mu-1}$ equals $\frac{d_\mu}{(\mu-1)!} \neq 0$. Using the above properties of the coefficients of the Taylor series of the function $\tilde{\psi}$ at zero, one easily proves that the germ of the function $\tilde{k}_1\tilde{k}_2$ at zero (and, thus, the germs of the functions $\tilde{k}_1^3$, $\tilde{k}_1^2 + \frac{2d_\mu}{(\mu-2)!}\tilde{k}_2^{\mu-2}$, and $\tilde{k}_2^{\mu-1}$) belongs to the Jacobian ideal $I_{\nabla\tilde{\psi}}$. It follows that the germs of the functions $\tilde{k}_1, \tilde{k}_1^2, \tilde{k}_2, \tilde{k}_2^2, \dots, \tilde{k}_2^{\mu-3}$ are generators of the local algebra $Q_{\nabla\tilde{\psi}}$ of the gradient map of the function $\tilde{\psi}$ at the origin (considered as a vector space), and that the differential operators $\frac{\partial}{\partial k_1}|_0, \frac{\partial}{\partial k_2}|_0, \frac{\partial^2}{\partial k_2^2}|_0, \dots, \frac{\partial^{\mu-3}}{\partial \tilde{k}_2^{\mu-3}}|_0, \frac{\partial^{\mu-2}}{\partial \tilde{k}_2^{\mu-2}}|_0 - d_\mu \frac{\partial^2}{\partial k_1^2}|_0$ are generators of its dual space $Q_{\nabla\tilde{\psi}}^*$. It follows from this, from the conditions on a deformation to be infinitesimally R^+ -versal, and from the versality theorem (see the theorems from [1, v. 1, 8.2 and 8.3]), that the deformation Ψ of the function ψ is R^+ -versal if and only if the vectors $\bar{d}_2, \dots, \bar{d}_{\mu-1}, \bar{d}_\mu - d_\mu \bar{d}_1 \in \mathbb{R}^l$ are linearly independent. Here $\bar{d}_i := \frac{\partial^{4-i}\tilde{\Psi}(0,0)}{\partial k_1^{3-i}\partial \lambda}$ for $i = 1, 2$, and $\bar{d}_i := \frac{\partial^{i-1}\tilde{\Psi}(0,0)}{\partial k_2^{i-2}\partial \lambda}$ for $i \geq 3$.

In order to compute the vectors $\bar{d}_\mu$, let us express the vectors $\bar{a}_{q,p} := \frac{\partial^{q+p+1}\Psi(0)}{\partial k_1^q \partial k_2^p \partial \lambda} = \frac{\partial^{q+p+1}F(0)}{\partial k_1^q \partial k_2^p \partial \lambda} + \dots$ in terms of the partial derivatives of the function F at the origin. Computations, similar to those for the vectors $\bar{a}_\mu = \frac{\partial^\mu F(0,0)}{\partial k_1^{\mu-1} \partial \lambda} + \dots \in \mathbb{R}^l$, see the proof of Statement 7.1, lead to the above formulae for $\bar{a}_{q,p}$, see the comment to Table (7.6). Performing the change of variables $(k_1, k_2) \rightarrow (\tilde{k}_1, \tilde{k}_2)$ and using the values of the coefficients λ_j from the proof of Statement 6.8, one obtains the required expressions for the vectors $\bar{d}_\mu$ in terms of the numbers $a_{q,p}$ and the vectors $\bar{a}_{q,p}$. This proves Statement 7.4.

Statement 7.5 is similarly proved on the basis of Theorem 4.6. Here one proves that the germ of the function $\tilde{k}_1^3\tilde{k}_2$ at the origin (and, thus, the germs of the even functions $\tilde{k}_1^6, \varepsilon\tilde{k}_1^4 + 2\eta\tilde{k}_1^2\tilde{k}_2^2, \eta\tilde{k}_1^2\tilde{k}_2^2 + \frac{\mu-3}{(2\mu-6)!}x_{e,\mu}^\varepsilon\tilde{k}_2^{2\mu-6}, \tilde{k}_1\tilde{k}_2^3$, and $\tilde{k}_2^{2\mu-4}$) belongs to the even Jacobian ideal $I_{\nabla\tilde{\psi}}^e$. It follows that the germs of the even functions $\tilde{k}_1^2, \tilde{k}_1\tilde{k}_2, \tilde{k}_2^2, \tilde{k}_1^4, \tilde{k}_2^4, \dots, \tilde{k}_2^{2\mu-8}$ are generators of the even local algebra $Q_{\nabla\tilde{\psi}}^e$ of the gradient map of the function $\tilde{\psi}$ at the origin (considered as a vector space), and that the differential operators $\frac{\partial^2}{\partial k_1^2}|_0, \frac{\partial^2}{\partial k_1\partial k_2}|_0, \frac{\partial^2}{\partial k_2^2}|_0, \frac{\partial^4}{\partial k_2^4}|_0, \dots, \frac{\partial^{2\mu-8}}{\partial \tilde{k}_2^{2\mu-8}}|_0, \frac{\partial^{2\mu-6}}{\partial \tilde{k}_2^{2\mu-6}}|_0 + \frac{\mu-3}{48}x_{e,\mu}^\varepsilon(\varepsilon\frac{\partial^4}{\partial k_1^4}|_0 - 12\eta\frac{\partial^4}{\partial k_1^2\partial k_2^2}|_0)$ are generators of its dual space $(Q_{\nabla\tilde{\psi}}^e)^*$. It follows from this and Theorem 4.6 that the deformation Ψ of the function ψ is R_O -versal if and only if the vectors $\bar{x}_{e,2}^\varepsilon, \dots, \bar{x}_{e,\mu-1}^\varepsilon, \bar{x}_{e,\mu}^\varepsilon + \frac{\mu-3}{48}x_{e,\mu}^\varepsilon\bar{x}_{e,1}^\varepsilon \in \mathbb{R}^l$ are linearly independent.

Statements 7.1, 7.2, 7.4, and 7.5 demonstrate that versal deformations are typical if the number l of parameters is large enough. More precisely: if $l \geq \mu - 1$ then the conditions from 7.1, 7.4 on a deformation of a germ f to be R^+ -versal (or the conditions from 7.2, 7.5 on an even deformation of an even germ f to be R_O -versal) are fulfilled for “typical” (resp. “typical even”) l -parameter deformations of f .

Here, by typical l -parameter (even) deformations of f we mean (even) deformations of f which form an open dense subspace in the space of all (even) l -parameter deformations of f in the convergence topology with a finite number of derivatives on each compact set.

References

- [1] V.I. Arnold, S.M. Gusein-Zade, A.N. Varchenko. *Singularities of Differentiable Maps*. Monographs in Mathematics, Birkhauser, Boston, 1985 and 1988.
- [2] V.I. Arnold, *Singularities of Caustics and Wave Fronts*. Kluwer Academic Publishers, 1990.
- [3] V.I. Arnol'd, V.A. Vasil'ev, V.V. Goryunov, O.V. Lyashko. *Singularities. Local and Global Theory*. Encyclopaedia of Math. Sci. **6**, Springer-Varlag, Berlin-Heidelberg-New York, 1993.

- [4] M. Beer, *Endliche Bestimmtheit und universelle Entfaltungen von Keimen mit Gruppenoperation*. Diplomarbeit, Regensburg 1976.
- [5] E.L. Lakshtanov, R.A. Minlos, *Spectrum of transfer-matrix two-particle bound states*, Func. Anal. i ego Pril. **38(3)** (2003), 52-69.
- [6] V.A. Malyshev, R.A. Minlos, *Linear Infinite-Particle Operators*. Amer. Math. Soc. Publ., 1995.
- [7] SH.S. Mamatov, R.A. Minlos, *Bound states of two-particle cluster operator*, Theor. and Math. Phys. **79(2)** (1989), 163-182.
- [8] R.A. Minlos, Ya.G. Sinay, *Study of the spectrum of stochastic operators appearing in the lattice models of gas*, Theor. and Math. Phys. **2** (1970), 230-243.
- [9] V. Poènaru, *Singularités C^∞ en présence de symétrie*, Lecture Notes in Math. **510**, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [10] P. Slodowy, *Einige Bemerkungen zur Entfaltung symmetrischer Funktionen*, Math. Z. **158** (1978), 157-170.
- [11] B.L. van der Waerden. *Algebra*. Springer-Verlag, Berlin-Heidelberg-New York, 1967.
- [12] G. Wassermann, *Classification of singularities with compact abelian symmetry*, Singularities, Banach Center Publications **20** (1988), 475-498.